arXiv:1509.00597v2 [math.AP] 10 Mar 2016

UNIQUENESS OF WEAK SOLUTIONS OF THE FULL COUPLED NAVIER-STOKES AND Q-TENSOR SYSTEM IN 2D

Francesco De Anna[1] and Arghir Zarnescu[2],[3]

[1]Université de Bordeaux
Institut de Mathématiques de Bordeaux
F-33405 Talence Cedex, France
Francesco.Deanna@math.u-bordeaux1.fr

[2]University of Sussex
Pevensey 3 5C17
Brighton BN1 9QF, UK
A.Zarnescu@sussex.ac.uk

[3] "Simion Stoilow" Institute of Mathematics
of the Romanian Academy
21 Calea Grivitei Street
010702 Bucharest, Romania

March 11, 2016

Abstract

This paper is devoted to the full system of incompressible liquid crystals, as modeled in the Q-tensor framework. The main purpose is to establish the uniqueness of weak solutions in a two dimensional setting, without imposing an extra regularity on the solutions themselves. This result only requires the initial data to fulfill the features which allow the existence of a weak solution. Thus, we also present a revisit of the global existence result in dimension two and three.

1. Introduction

The main aim of this article is to prove the uniqueness of weak solutions for a type of coupled Navier-Stokes and Q-tensor systems proposed in [5] and studied numerically and analytically in [1,14,16–18,36]. This type of system models nematic liquid crystals and provides in a certain sense an extension of the classical Ericksen-Leslie model [14], whose uniqueness of weak solutions was proved in [38]. In the remainder of this introduction we will briefly present the equations and state our main result.

The system models the evolution of liquid crystal molecules together with the underlying flow, through a parabolic-type system coupling an incompressible Navier-Stokes system with a nonlinear convection-diffusion system. The local orientation of the molecules is described through a function Q taking values from $\mathbb{R}_+ \times \Omega \subset \mathbb{R}_+ \times \mathbb{R}^d$,$d = 2, 3$ into the set of so-called d-dimensional Q-tensors that is

$$S_0^{(d)} \overset{\text{def}}{=} \left\{ Q \in \mathbb{M}^{d\times d}; Q_{ij} = Q_{ji}, \mathrm{tr}(Q) = 0, i, j = 1, \dots, d \right\}$$

(the most relevant physical situations being $d = 2, 3$). The evolution of the Q's is driven by a gradient flow of the free energy of the molecules as well as the transport, distortion and alignment effects caused by the flow. The flow field $u : \mathbb{R}_+ \times \Omega \to \mathbb{R}^d$ satisfies a forced incompressible Navier-Stokes system, with the forcing provided by the additional, non-Newtonian stress caused by the molecules orientations, thus expressed in terms of Q. We restrict ourselves to the case $\Omega = \mathbb{R}^d$ and work with non-dimensional quantities. The evolution of Q is given by:

$$\partial_t Q + u \cdot \nabla Q - S(\nabla u, Q) = -\Gamma \frac{\partial \mathcal{F}_e}{\partial Q} \tag{1}$$

with $\Gamma > 0$. Here

$$\mathcal{F}_e(Q) = \int_{\mathbb{R}^d} \frac{L}{2}|\nabla Q|^2 + \left(\frac{a}{2}\mathrm{tr}(Q^2) - \frac{b}{3}\mathrm{tr}(Q^3) + \frac{c}{4}\mathrm{tr}^2(Q^2)\right) dx \tag{2}$$

is the free energy of the liquid crystal molecules and $\frac{\partial \mathcal{F}_e}{\partial Q}$ denotes the variational derivative. The L, a, b, c constants are specific to the material with:

$$L > 0 \text{ and } a, b, c \in \mathbb{R},\ c > 0 \tag{3}$$

If $u = 0$ the Q-tensor equation would simply be a gradient flow of the free energy. For $u \neq 0$ the molecules are transported by the flow (as indicated by the convective derivative $\partial_t + u \cdot \nabla$) as well as being tumbled and aligned by the flow, fact described by the term

$$S(\nabla u, Q) \overset{\text{def}}{=} (\xi D + \Omega)(Q + \frac{1}{d}Id) + (Q + \frac{1}{d}Id)(\xi D - \Omega) - 2\xi(Q + \frac{1}{d}Id)\text{tr}(Q\nabla u) \tag{4}$$

where $D \overset{def}{=} \frac{1}{2}\big(\nabla u + (\nabla u)^T\big)$ and $\Omega \overset{def}{=} \frac{1}{2}\big(\nabla u - (\nabla u)^T\big)$ are, respectively, the symmetric part and the antisymmetric part, of the velocity gradient matrix ∇u. The constant ξ is specific to the liquid crystal material.

The flow satisfies the forced Navier-Stokes system:

$$\begin{aligned} \partial_t u + (u \cdot \nabla)u &= \nu\Delta u + \nabla p + \lambda \nabla \cdot (\tau + \sigma) \\ \nabla \cdot u &= 0 \end{aligned}$$

where $\nu, \lambda > 0$ with λ measuring the ratio of the elastic effects (produced by the liquid crystal molecules) to that of the diffusive effects. The forcing is provided by the additional stress caused by the presence of the liquid crystal molecules, more specifically we have the symmetric part of the additional stress tensor:

$$\tau \overset{\text{def}}{=} \Big[-\xi\Big(Q + \frac{1}{d}Id\Big)H - \xi H\big(Q + \frac{1}{d}Id\big) + 2\xi\big(Q + \frac{1}{d}Id\big)QH - L\nabla Q \odot \nabla Q\Big] \tag{5}$$

and the antisymmetric part:

$$\sigma \overset{\text{def}}{=} QH - HQ \tag{6}$$

where we denoted

$$H \overset{\text{def}}{=} -\frac{\partial \mathcal{F}_e}{\partial Q} = L\Delta Q - aQ + b[Q^2 - \frac{\text{tr}(Q^2)}{d}Id] - cQ\text{tr}(Q^2) \tag{7}$$

Summarising we have the coupled system:

$$\begin{aligned} (\partial_t + u \cdot \nabla)Q - S(\nabla u, Q) =& \Gamma(L\Delta Q - aQ + b[Q^2 - \frac{\text{tr}(Q^2)}{d}Id] - cQ\text{tr}(Q^2)) \\ \partial_t u + (u \cdot \nabla)u =& \nu\Delta u + \nabla p + \lambda\nabla \cdot (QH - HQ) \\ &+ \lambda\nabla \cdot \Big[-\xi\Big(Q + \frac{1}{d}Id\Big)H - \xi H\big(Q + \frac{1}{d}Id\big) + 2\xi\big(Q + \frac{1}{d}Id\big)QH - L\nabla Q \odot \nabla Q\Big] \\ \nabla \cdot u =& 0 \end{aligned} \tag{8}$$

where $\Gamma, L, \nu, c > 0$, $a, b \in \mathbb{R}$. Let us observe that this is a slight extension of the system considered in [36], where $\lambda = 1$. However, this does not create any major difficulties compared to equations in [36] but it is more relevant from a physical point of view.

The main result of the paper is the uniqueness of weak solutions, which are defined in a rather standard manner:

Definition of weak solutions *A pair (Q, u) is called a weak solution of the system (8), subject to initial data*

$$Q(0, x) = \bar{Q}(x) \in H^1(\mathbb{R}^d; S_0^{(d)}),\ u(0, x) = \bar{u}(x) \in L^2(\mathbb{R}^d; \mathbb{R}^d), \nabla \cdot \bar{u} = 0 \ \textit{ in } \mathcal{D}'(\mathbb{R}^d) \tag{9}$$

if $Q \in L^\infty_{loc}(\mathbb{R}_+; H^1) \cap L^2_{loc}(\mathbb{R}_+; H^2)$, $u \in L^\infty_{loc}(\mathbb{R}_+; L^2) \cap L^2_{loc}(\mathbb{R}_+; H^1)$ and for every compactly supported $\varphi \in C^\infty([0, \infty) \times \mathbb{R}^d; S_0^{(d)})$, $\psi \in C^\infty([0, \infty) \times \mathbb{R}^d; \mathbb{R}^d)$ with $\nabla \cdot \psi = 0$ we have

$$\begin{aligned} &\int_0^\infty \int_{\mathbb{R}^d} (-Q \cdot \partial_t \varphi - \Gamma L \Delta Q \cdot \varphi) - Q \cdot u \nabla_x \varphi dx\, dt \\ -&\int_0^\infty \int_{\mathbb{R}^d} (\xi D + \Omega)(Q + \frac{1}{d}Id) \cdot \varphi + (Q + \frac{1}{d}Id)(\xi D - \Omega) \cdot \varphi - 2\xi(Q + \frac{1}{d}Id)tr(Q\nabla u) \cdot \varphi\, dx\, dt \\ &= \int_{\mathbb{R}^d} \bar{Q}(x) \cdot \varphi(0, x)\, dx + \Gamma \int_0^\infty \int_{\mathbb{R}^d} \Big\{ -aQ + b[Q^2 - \frac{tr(Q^2)}{d}Id] - cQtr(Q^2)\Big\} \cdot \varphi\ dx\, dt \end{aligned} \tag{10}$$

and

$$\int_0^\infty \int_{\mathbb{R}^d} -u\partial_t \psi - u_\alpha u_\beta \partial_\alpha \psi_\beta + \nu \nabla u \nabla \psi\ dt\, dx - \int_{\mathbb{R}^d} \bar{u}(x)\psi(0, x)\, dx$$

$$= L\lambda \int_0^\infty \int_{\mathbb{R}^d} Q_{\gamma\delta,\alpha} Q_{\gamma\delta,\beta} \psi_{\alpha,\beta} - Q_{\alpha\gamma} \Delta Q_{\gamma\beta} \psi_{\alpha,\beta} + \Delta Q_{\alpha\gamma} Q_{\gamma\beta} \psi_{\alpha,\beta} \, dx\, dt$$

$$+\xi\lambda \int_0^\infty \int_{\mathbb{R}^d} \left(Q_{\alpha\gamma} + \frac{\delta_{\alpha\gamma}}{d}\right) H_{\gamma\beta} \psi_{\alpha,\beta} + H_{\alpha\gamma} \left(Q_{\gamma\beta} + \frac{\delta_{\gamma\beta}}{d}\right) \psi_{\alpha,\beta} - 2(Q_{\alpha\beta} + \frac{\delta_{\alpha\beta}}{d}) Q_{\gamma\delta} H_{\gamma\delta} \psi_{\alpha,\beta} \, dx\, dt \qquad (11)$$

We can now state our main result, which is the existence and uniqueness of weak solutions:

Theorem 1.1. *Let $d = 2, 3$ and take*

$$Q(0,x) = \bar{Q}(x) \in H^1(\mathbb{R}^d; S_0^{(d)}),\ u(0,x) = \bar{u}(x) \in L^2(\mathbb{R}^d), \nabla \cdot \bar{u} = 0 \ \text{ in } \mathcal{D}'(\mathbb{R}^d)$$

Then the system (8) *admits a global weak solution. Moreover if $d = 2$, then uniqueness holds.*

Remark 1.2. *With minor modifications to the proof, that are left to the interested reader, the result also holds when the system is $d = 2$ in the domain but $d = 3$ in the target, which physically corresponds to a situation where there is no dependence in one of the three spatial directions.*

The main part of the theorem is about uniqueness, as the existence part is just a fairly straightforward revisit of the arguments in [36]. The main difficulties associated with treating the system (8) are related to the presence of the Navier-Stokes part. One can essentially think of the system as a highly non-trivial perturbation of a Navier-Stokes system. It is known that for Navier-Stokes alone the uniqueness of weak solutions in $2D$ can be achieved through rather standard arguments, while in $3D$ it is a major open problem.

The extended system that we deal with has an intermediary position, as the perturbation produced by the presence of the additional stress-tensor generates significant technical difficulties related in the first place to the weak norms available for the u term. A rather common way of dealing with this issue is by using a weak norm for estimating the difference between the two weak solutions, a norm that is below the natural spaces in which the weak solutions are defined. This approach was used before in the context of the related Leslie-Ericksen model [20] as well as for the usual Navier-Stokes system in [15] and [29].

In our case, for technical convenience we use a homogeneous Sobolev space, namely $\dot{H}^{-\frac{1}{2}}$. The fact that the initial data for the difference is zero (i.e. $(\delta u, \delta Q)_{t=0} = 0$) helps in controlling the difference in such a low regularity space. However, one of the main reasons for chosing the homogeneous setting is a specific product law, see Theorem A.1 in the Appendix. The mentioned theorem shows that the product is a bounded operator from $\dot{H}^s(\mathbb{R}^2) \times \dot{H}^t(\mathbb{R}^2)$ into $\dot{H}^{s+t-1}(\mathbb{R}^2)$, for any $|s|$, $|t| \le 1$ such that $s + t$ is positive. We note that evaluating the difference at regularity level $s = 0$ i.e. in L^2, would only allow to prove a weak-strong uniqueness result, along the lines of [35]. Working in a negative Sobolev space, $\dot{H}^s$ with $s \in (-1, 0)$ allows to capture the uniqueness of weak solutions. We expect that a similar proof would work in any $\dot{H}^s$ with $s \in (-1, 0)$ and our choice $s = -\frac{1}{2}$ is just for convenience.

Our main work is to obtain the delicate double-logarithmic type estimates that lead to an Osgood lemma, a generalization of the Gronwall inequality (see [2], Lemma 3.4). Indeed the uniqueness reduces to an estimate of the following type:

$$\Phi'(t) \le \chi(t) \Big\{ \Phi(t) + \Phi(t) \ln\Big(1 + e + \frac{1}{\Phi(t)}\Big) + \Phi(t) \ln\Big(1 + e + \frac{1}{\Phi(t)}\Big) \ln\ln\Big(1 + e + \frac{1}{\Phi(t)}\Big) \Big\},$$

where $\Phi(t)$ is a norm of the difference between two solutions and χ is apriori in L^1_{loc}.

In addition to these there are some difficulties that are specific to this system. These are of two different types, being related to:

- controlling the “extraneous” maximal derivatives: that is the highest derivatives in u that appear in the Q equation and the highest derivatives in Q that appear in the u equation,
- controlling the high powers of Q , such as $Q\mathrm{tr}(Q^2)$ in particular those that interact with u terms (such as $Q\mathrm{tr}(Q\nabla u)$).

The first difficulty is dealt with by taking into account the specific feature of the coupling that allows for the *cancellation of the worst terms*, when considering certain physically meaningful combinations of terms. This feature is explored in the next section where we revisit and revise the existence proof from [36]. In what concerns the second difficulty, this is overcome by delicate harmonic analysis arguments leading to the double logarithmic estimates mentioned before.

The paper is organised as follows: in the next section we revisit the existence arguments done in cite [36], providing a slight adaptation to our case and a minor correction to one of the estimates used there. The main benefit of this section is that it exhibits in a simple setting a number of cancellations that are later-on crucial for the uniqueness argument. In the third section we start by introducing a number of technical harmonic analysis tools related to the Littlewood-Paley theory and then use them in the proof of our main result. Some standard but perhaps less-known tools, toghether with some more technical estimates are postponed into the appendices.

Notations and conventions. Let $S_0^{(d)} \subset \mathbb{M}^{d\times d}$ denote the space of Q-tensors in dimension d, i.e.

$$S_0^{(d)} \overset{\text{def}}{=} \big\{Q \in \mathbb{M}^{d\times d}; Q_{ij} = Q_{ji}, \mathrm{tr}(Q) = 0, i, j = 1, \dots, d\big\}$$

We use the Einstein summation convention, that is we assume summation over repeated indices.

We define the Frobenius norm of a matrix $|Q| \overset{\text{def}}{=} \sqrt{\mathrm{tr}Q^2} = \sqrt{Q_{\alpha\beta}Q_{\alpha\beta}}$ and define Sobolev spaces of Q-tensors in terms of this norm. For instance $H^1(\mathbb{R}^d, S_0^{(d)}) \overset{\text{def}}{=} \{Q : \mathbb{R}^d \to S_0^{(d)}, \int_{\mathbb{R}^d} |\nabla Q(x)|^2 + |Q(x)|^2\, dx < \infty\}$ where $|\nabla Q|^2(x) \overset{\text{def}}{=} Q_{\alpha\beta,\gamma}(x)Q_{\alpha\beta,\gamma}(x)$ with $Q_{\alpha\beta,\gamma} \overset{\text{def}}{=} \partial_\gamma Q_{\alpha\beta}$. For $A, B \in S_0^{(d)}$ we denote $A : B = \mathrm{tr}(AB)$, $|A| = \sqrt{\mathrm{tr}(A^2)}$ and $\|(A,\, B)\|_X = \|A\|_X + \|B\|_X$, for any suitable Banach space X. We also denote $\Omega_{\alpha\beta} \overset{\text{def}}{=} \frac{1}{2}(\partial_\beta u_\alpha - \partial_\alpha u_\beta)$,$u_{\alpha,\beta} \overset{\text{def}}{=} \partial_\beta u_\alpha$ and $(\nabla Q \odot \nabla Q)_{ij} = Q_{\alpha\beta,i}Q_{\alpha\beta,j}$.

2. The energy decay, apriori estimates and scaling

In the absence of the flow, when $u = 0$ in the equations (8), the free energy is a Lyapunov functional of the system. If $u \neq 0$ we still have a Lyapunov functional for (8) but this time one that includes the kinetic energy of the system. These estimates provide as usually the basis for obtaining apriori estimates for the system. The propositions in this section show this and their proofs follow closely the ones of the similar propositions in [36] where they were done for the case $\lambda = 1$. The reason for including them is to display in relatively simple setting the cancellations that will appear again in the proof of the uniqueness theorem but in a much more complicated framework. We have:

Proposition 2.1. *The system* (8) *has a Lyapunov functional:*

$$E(t) \overset{\text{def}}{=} \frac{1}{2}\int_{\mathbb{R}^d} |u|^2(t,x)\,dx + \int_{\mathbb{R}^d} \frac{L\lambda}{2}|\nabla Q|^2(t,x) + \lambda\Big(\frac{a}{2}tr(Q^2(t,x)) - \frac{b}{3}tr(Q^3(t,x)) + \frac{c}{4}tr^2(Q^2(t,x))\Big)\,dx \quad (12)$$

If $d = 2, 3$ and (Q, u) is a smooth solution of (8) such that $Q \in L^\infty(0,T;H^1(\mathbb{R}^d)) \cap L^2(0,T;H^2(\mathbb{R}^d))$ and $u \in L^\infty(0,T;L^2(\mathbb{R}^d)) \cap L^2(0,T;H^1(\mathbb{R}^d))$ then, for all $t < T$, we have:

$$\frac{d}{dt}E(t) = -\nu\int_{\mathbb{R}^d} |\nabla u|^2\,dx - \Gamma\lambda\int_{\mathbb{R}^d} tr\left(L\Delta Q - aQ + b[Q^2 - \frac{tr(Q^2)}{d}Id] - cQtr(Q^2)\right)^2 dx \le 0 \quad (13)$$

Proof. We multiply the first equation in (8) to the right by $-\lambda H$, take the trace, integrate over $\mathbb{R}^d$ and by parts and sum with the second equation multiplied by u and integrated over $\mathbb{R}^d$ and by parts (let us observe that because of our assumptions on Q and u we do not have boundary terms, when integrating by parts). We obtain:

$$\frac{d}{dt}\int_{\mathbb{R}^d} \frac{1}{2}|u|^2 + \frac{L\lambda}{2}|\nabla Q|^2 + \lambda\Big(\frac{a}{2}\mathrm{tr}(Q^2) - \frac{b}{3}\mathrm{tr}(Q^3) + \frac{c}{4}\mathrm{tr}^2(Q^2)\Big)\,dx$$

$$+\nu\int_{\mathbb{R}^d} |\nabla u|^2\,dx + \Gamma\lambda\int_{\mathbb{R}^d} \mathrm{tr}\left(L\Delta QL - aQ + b[Q^2 - \frac{\mathrm{tr}(Q^2)}{d}Id] - cQ\mathrm{tr}(Q^2)\right)^2 dx$$

$$= \underbrace{\lambda\int_{\mathbb{R}^d} u\cdot\nabla Q_{\alpha\beta}\left(-aQ_{\alpha\beta} + b[Q_{\alpha\gamma}Q_{\gamma\beta} - \frac{\delta_{\alpha\beta}}{d}\mathrm{tr}(Q^2)] - cQ_{\alpha\beta}\mathrm{tr}(Q^2))\right)dx}_{\overset{\text{def}}{=}\mathcal{I}}$$

$$\underbrace{+\lambda\int_{\mathbb{R}^d}(-\Omega_{\alpha\gamma}Q_{\gamma\beta} + Q_{\alpha\gamma}\Omega_{\gamma\beta})\left(-aQ_{\alpha\beta} + b[Q_{\alpha\delta}Q_{\delta\beta} - \frac{\delta_{\alpha\beta}}{d}\mathrm{tr}(Q^2)] - cQ_{\alpha\beta}\mathrm{tr}(Q^2))\right)dx}_{\overset{\text{def}}{=}\mathcal{II}}$$

$$-\lambda\xi\underbrace{\int_{\mathbb{R}^d}(Q_{\alpha\gamma} + \frac{\delta_{\alpha\gamma}}{d})D_{\gamma\beta}H_{\alpha\beta}dx}_{\overset{\text{def}}{=}\mathcal{J}_1} - \lambda\xi\underbrace{\int_{\mathbb{R}^d}D_{\alpha\gamma}(Q_{\gamma\beta} + \frac{\delta_{\gamma\beta}}{d})H_{\alpha\beta}dx}_{\overset{\text{def}}{=}\mathcal{J}_2} + 2\lambda\xi\underbrace{\int_{\mathbb{R}^d}(Q_{\alpha\beta} + \frac{\delta_{\alpha\beta}}{d})H_{\alpha\beta}\mathrm{tr}(Q\nabla u)dx}_{\overset{\text{def}}{=}\mathcal{J}_3}$$

$$+L\lambda\underbrace{\int_{\mathbb{R}^d}u_\gamma Q_{\alpha\beta,\gamma}\Delta Q_{\alpha\beta}\,dx}_{\overset{\text{def}}{=}\mathcal{A}}\underbrace{-\frac{L\lambda}{2}\int_{\mathbb{R}^d}u_{\alpha,\gamma}Q_{\gamma\beta}\Delta Q_{\alpha\beta}\,dx}_{\overset{\text{def}}{=}\mathcal{B}}$$

$$\underbrace{+\frac{L\lambda}{2}\int_{\mathbb{R}^d} u_{\gamma,\alpha}Q_{\gamma\beta}\Delta Q_{\alpha\beta}\,dx}_{\overset{\text{def}}{=}\mathcal{C}}\underbrace{+\frac{L\lambda}{2}\int_{\mathbb{R}^d} Q_{\alpha\gamma}u_{\gamma,\beta}\Delta Q_{\alpha\beta}\,dx}_{\mathcal{C}}\underbrace{-\frac{L\lambda}{2}\int_{\mathbb{R}^d} Q_{\alpha\gamma}u_{\beta,\gamma}\Delta Q_{\alpha\beta}\,dx}_{\mathcal{B}}$$

$$+L\lambda\underbrace{\int_{\mathbb{R}^d} Q_{\gamma\delta,\alpha}Q_{\gamma\delta,\beta}u_{\alpha,\beta}\,dx}_{\overset{\text{def}}{=}\mathcal{AA}}\underbrace{-L\lambda\int_{\mathbb{R}^d} Q_{\alpha\gamma}\Delta Q_{\gamma\beta}u_{\alpha,\beta}\,dx}_{\overset{\text{def}}{=}\mathcal{CC}}\underbrace{+L\lambda\int_{\mathbb{R}^d} \Delta Q_{\alpha\gamma}Q_{\gamma\beta}u_{\alpha,\beta}\,dx}_{\overset{\text{def}}{=}\mathcal{BB}}$$

$$+\lambda\xi\underbrace{\int_{\mathbb{R}^d}\big(Q_{\alpha\gamma}+\frac{\delta_{\alpha\gamma}}{d}\big)H_{\gamma\beta}u_{\alpha,\beta}\,dx}_{\overset{\text{def}}{=}\mathcal{JJ}_1}+\lambda\xi\underbrace{\int_{\mathbb{R}^d}H_{\alpha\gamma}\big(Q_{\gamma\beta}+\frac{\delta_{\gamma\beta}}{d}\big)u_{\alpha,\beta}\,dx}_{\overset{\text{def}}{=}\mathcal{JJ}_2}-2\lambda\xi\underbrace{\int_{\mathbb{R}^d}\big(Q_{\alpha\beta}+\frac{\delta_{\alpha\beta}}{d}\big)u_{\alpha,\beta}\mathrm{tr}(QH)\,dx}_{\overset{\text{def}}{=}\mathcal{JJ}_3}$$

$$=\underbrace{-L\lambda\int_{\mathbb{R}^d} u_{\alpha,\gamma}Q_{\gamma\beta}\Delta Q_{\alpha\beta}\,dx}_{2\mathcal{B}}\underbrace{+L\lambda\int_{\mathbb{R}^d} u_{\gamma,\alpha}Q_{\gamma\beta}\Delta Q_{\alpha\beta}\,dx}_{2\mathcal{C}}$$

$$\underbrace{-L\lambda\int_{\mathbb{R}^d} Q_{\alpha\gamma}\Delta Q_{\gamma\beta}u_{\alpha,\beta}\,dx}_{\mathcal{CC}}\underbrace{+L\lambda\int_{\mathbb{R}^d} \Delta Q_{\alpha\gamma}Q_{\gamma\beta}u_{\alpha,\beta}\,dx}_{\mathcal{BB}}=0 \tag{14}$$

where $\mathcal{I}=0$ (since $\nabla\cdot u=0$), $\mathcal{II}=0$ (since $Q_{\alpha\beta}=Q_{\beta\alpha}$) and for the second equality we used

$$\underbrace{\int_{\mathbb{R}^d} u_\gamma Q_{\alpha\beta,\gamma}\Delta Q_{\alpha\beta}\,dx}_{\mathcal{A}}+\underbrace{\int_{\mathbb{R}^d} Q_{\gamma\delta,\alpha}Q_{\gamma\delta,\beta}u_{\alpha,\beta}\,dx}_{\mathcal{AA}}=\int_{\mathbb{R}^d} u_\gamma Q_{\alpha\beta,\gamma}\Delta Q_{\alpha\beta}\,dx$$

$$-\int_{\mathbb{R}^d} Q_{\gamma\delta,\alpha}Q_{\gamma\delta,\beta\beta}u_\alpha\,dx-\int_{\mathbb{R}^d} Q_{\gamma\delta,\alpha\beta}Q_{\gamma\delta,\beta}u_\alpha\,dx=\int_{\mathbb{R}^d}\frac{1}{2}Q_{\gamma\delta,\beta}Q_{\gamma\delta,\beta}u_{\alpha,\alpha}\,dx=0$$

together with $Q_{\alpha\alpha}=H_{\alpha\alpha}=u_{\alpha,\alpha}=0$, $\mathcal{J}_3=\mathcal{JJ}_3$ and

$$\mathcal{J}_1+\mathcal{J}_2=\int_{\mathbb{R}^d}\frac{1}{2}Q_{\alpha\gamma}u_{\gamma,\beta}H_{\alpha\beta}+\frac{1}{2}Q_{\alpha\gamma}u_{\beta,\gamma}H_{\alpha\beta}+\frac{1}{2}u_{\alpha,\gamma}Q_{\gamma\beta}H_{\alpha\beta}+\frac{1}{2}u_{\gamma,\alpha}Q_{\gamma\beta}H_{\alpha\beta}\,dx$$

$$+\frac{2}{d}\int_{\mathbb{R}^d}D_{\alpha\beta}H_{\alpha\beta}=\int_{\mathbb{R}^d}\frac{1}{2}\big(Q_{\alpha\gamma}u_{\gamma,\beta}H_{\alpha\beta}+u_{\gamma,\alpha}Q_{\gamma\beta}H_{\alpha\beta}\big)+\frac{1}{2}\big(Q_{\alpha\gamma}u_{\beta,\gamma}H_{\alpha\beta}+u_{\alpha,\gamma}Q_{\gamma\beta}H_{\alpha\beta}\big)\,dx$$

$$+\frac{1}{d}\int_{\mathbb{R}^d}(u_{\alpha,\beta}+u_{\beta,\alpha})H_{\alpha\beta}\,dx=\int_{\mathbb{R}^d}H_{\beta\alpha}Q_{\alpha\gamma}u_{\gamma,\beta}+Q_{\gamma\alpha}H_{\alpha\beta}u_{\beta,\gamma}\,dx+\frac{2}{d}\int_{\mathbb{R}^d}u_{\alpha,\beta}H_{\alpha\beta}\,dx=\mathcal{JJ}_1+\mathcal{JJ}_2.$$

Finally, the last equality in (14) is a consequence of the straightforward identities $2\mathcal{B}+\mathcal{BB}=2\mathcal{C}+\mathcal{CC}=0$.
☐

It can be easily checked that the system has a scaling, namely we have:

Lemma 2.2. *Let (Q,u,p) be a solution of (8). Then letting*

$$u_\delta\overset{\text{def}}{=}\delta u(\delta x,\delta^2 t),\ Q_\delta\overset{\text{def}}{=}Q(\delta x,\delta^2 t),\ p_\delta(x,t)\overset{\text{def}}{=}\delta^2 p(\delta x,\delta^2) \tag{15}$$

we have that $(Q_\delta,u_\delta,p_\delta)$ satisfy (8) with $F(Q)=-aQ+b[Q^2-\frac{tr(Q^2)}{3}Id]-cQtr(Q^2)$ replaced by $F_\delta(Q_\delta)=\delta^2\left[-aQ_\delta+b[(Q_\delta)^2-\frac{tr(Q_\delta^2)}{3}-cQ_\delta tr(Q_\delta)^2\right]$. We note that, in dimension two, the space $\dot H^1(\mathbb{R}^2)\times L^2(\mathbb{R}^2)$ is invariant by the scaling.

In the following we assume that there exists a smooth solution of (8) and obtain estimates on the behaviour of various norms.

Proposition 2.3. *Let (Q,u) be a smooth solution of (8) in dimension $d=2$ or $d=3$, with restriction (3), and smooth initial data $(\bar Q(x),\bar u(x))$, that decays fast enough at infinity so that we can integrate by parts in space (for any $t\geq 0$) without boundary terms. We assume that $|\xi|<\xi_0$ where ξ_0 is an explicitly computable constant, scale invariant, depending on $a,b,c,d,\Gamma,\nu,\lambda$.*

For $(\bar Q,\bar u)\in H^1\times L^2_x$,we have

$$\|Q(t,\cdot)\|_{H^1}\leq C_1+\bar C_1 e^{\bar C_1 t}\|\bar Q\|_{H^1},\forall t\geq 0 \tag{16}$$

with $C_1,\bar C_1$ depending on $(a,b,c,d,\Gamma,L,\nu,\bar Q,\bar u)$. Moreover

$$\|u(t,\cdot)\|^2_{L^2_x}+\nu\int_0^t\|\nabla u\|^2_{L^2_x}\leq C_1 \tag{17}$$

Proof. We denote:

$$X_{\alpha\beta} \overset{\mathrm{def}}{=} L\Delta Q_{\alpha\beta} - cQ_{\alpha\beta}\mathrm{tr}(Q^2),\ \alpha,\beta = 1,2,3 \tag{18}$$

Then equation (13) becomes:

$$\frac{d}{dt}E(t) + \nu\|\nabla u\|^2_{L^2_x} + \Gamma\lambda L^2\|\Delta Q\|^2_{L^2_x} + \Gamma\lambda c^2\|Q\|^6_{L^6} - 2cL\Gamma\lambda\int_{\mathbb{R}^d}\Delta Q_{\alpha\beta}Q_{\alpha\beta}\mathrm{tr}(Q^2)\,dx + a^2\Gamma\lambda\|Q\|^2_{L^2_x} +$$
$$+ b^2\Gamma\lambda\int_{\mathbb{R}^d}\mathrm{tr}\left(Q^2 - \frac{\mathrm{tr}(Q^2)}{d}\right)^2 dx \le 2a\Gamma\lambda\underbrace{\int_{\mathbb{R}^d}\mathrm{tr}(XQ)\,dx}_{\overset{\mathrm{def}}{=}\mathcal{I}} - 2b\Gamma\lambda\underbrace{\int_{\mathbb{R}^d}\mathrm{tr}(XQ^2)\,dx}_{\overset{\mathrm{def}}{=}\mathcal{J}} + 2ab\Gamma\lambda\int_{\mathbb{R}^d}\mathrm{tr}(Q^3)\,dx \tag{19}$$

Integrating by parts we have:

$$-2cL\Gamma\lambda\int_{\mathbb{R}^d}\Delta Q_{\alpha\beta}Q_{\alpha\beta}\mathrm{tr}(Q^2)dx = 2cL\Gamma\lambda\int_{\mathbb{R}^d}Q_{\alpha\beta,k}Q_{\alpha\beta,k}\mathrm{tr}(Q^2)dx + 2cL\Gamma\lambda\int_{\mathbb{R}^d}Q_{\alpha\beta,k}Q_{\alpha\beta}\partial_k\left(\mathrm{tr}(Q^2)\right)dx$$
$$= 2cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla Q|^2\mathrm{tr}(Q^2)\,dx + cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla\left(\mathrm{tr}(Q^2)\right)|^2\,dx \ge 0 \tag{20}$$

(where for the last inequality we used the assumption (3) and $L,\Gamma,\lambda > 0$). One can easily see that

$$\mathcal{I} = -\frac{L}{2}\|\nabla Q\|^2_{L^2_x} - c\|Q\|^4_{L^4} \tag{21}$$

On the other hand, for any $\varepsilon > 0$ and $\tilde{C} = \tilde{C}(\varepsilon, c)$ an explicitly computable constant, we have:

$$\mathcal{J} = L\int_{\mathbb{R}^d}Q_{\alpha\beta,kk}Q_{\alpha\gamma}Q_{\gamma\beta}\,dx - c\int_{\mathbb{R}^d}\mathrm{tr}(Q^2)\mathrm{tr}(Q^3)\,dx$$
$$\le -L\int_{\mathbb{R}^d}Q_{\alpha\beta,k}Q_{\alpha\gamma,k}Q_{\gamma\beta}\,dx - L\int_{\mathbb{R}^d}Q_{\alpha\beta,k}Q_{\alpha\gamma}Q_{\gamma\beta,k} + \int_{\mathbb{R}^d}\mathrm{tr}(Q^2)\left(\frac{\tilde{C}}{\varepsilon}\mathrm{tr}(Q^2) + \varepsilon\mathrm{tr}^2(Q^2)\right)dx$$
$$\le L\varepsilon\int_{\mathbb{R}^d}|\nabla Q|^2\mathrm{tr}(Q^2)\,dx + \frac{\tilde{C}}{\varepsilon}\|\nabla Q\|^2_{L^2_x} + \int_{\mathbb{R}^d}\mathrm{tr}(Q^2)\left(\frac{\tilde{C}}{\varepsilon}\mathrm{tr}(Q^2) + \varepsilon\mathrm{tr}^2(Q^2)\right)dx$$

Using the last three relations in (19) we obtain:

$$\frac{d}{dt}E(t) + \nu\|\nabla u\|^2_{L^2_x} + \Gamma\lambda L^2\|\Delta Q\|^2_{L^2_x} + c^2\Gamma\lambda\|Q\|^6_{L^6} + a^2\Gamma\lambda\|Q\|^2_{L^2_x} + 2cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla Q|^2\mathrm{tr}(Q^2)\,dx$$
$$+ cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla\left(\mathrm{tr}(Q^2)\right)|^2\,dx \le 2|a|\Gamma\lambda\Big(\frac{L}{2}\|\nabla Q\|^2_{L^2_x} + c\|Q\|^4_{L^4}\Big) + 2|b|\Gamma\lambda L\varepsilon\int_{\mathbb{R}^d}|\nabla Q|^2\mathrm{tr}(Q^2)\,dx$$
$$+ 2|b|\Gamma\lambda\frac{\tilde{C}}{\varepsilon}\|\nabla Q\|^2_{L^2_x} + 2|b|\Gamma\lambda\int_{\mathbb{R}^d}\mathrm{tr}(Q^2)\left(\frac{\tilde{C}}{\varepsilon}\mathrm{tr}(Q^2) + \varepsilon\mathrm{tr}^2(Q^2)\right)dx + 2|ab|\Gamma\lambda\Big(\varepsilon\|Q\|^2_{L^2_x} + \frac{\tilde{C}}{\varepsilon}\|Q\|^4_{L^4}\Big)$$

Taking ε small enough we can absorb all the terms with an epsilon coefficient on the right into the left hand side, and we are left with

$$\frac{d}{dt}E(t) + \nu\|\nabla u\|^2_{L^2_x} + \Gamma\lambda L^2\|\Delta Q\|^2_{L^2_x} + \Gamma\lambda c^2\|Q\|^6_{L^6} + \Gamma\lambda a^2\|Q\|^2_{L^2_x}$$
$$+ 2cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla Q|^2\mathrm{tr}(Q^2)\,dx + cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla\left(\mathrm{tr}(Q^2)\right)|^2\,dx \le \bar{C}\left(\|\nabla Q\|^2_{L^2_x} + \|Q\|^4_{L^4}\right) \tag{22}$$

with $\bar{C} = \bar{C}(a,b,c)$.

The last relation is not yet enough because the Q terms without derivatives in $E(t)$ are not summing to a positive number. However, let us note that, if $a > 0$ we obtain the a-priori estimates by using the inequality $\mathrm{tr}(Q^3) \le \frac{3}{8}\mathrm{tr}(Q^2) + \mathrm{tr}(Q^2)^2$. If $a \le 0$ we have to estimate separately $\|Q\|_{L^2_x}$ and this ask for a smallness condition for ξ.

We need to control in some sense low frequencies of Q. To this end we multiply the first equation in (8) by Q, take the trace, integrate over $\mathbb{R}^d$ and by parts and we obtain:

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^d}|Q|^2(t,x)\,dx = \Gamma\Big(-L\int_{\mathbb{R}^d}|\nabla Q|^2\,dx - a\int_{\mathbb{R}^d}|Q(x)|^2\,dx + b\int_{\mathbb{R}^d}\mathrm{tr}(Q^3)\,dx - c\int_{\mathbb{R}^d}|Q|^4\,dx\Big)$$
$$+ \underbrace{\int_{\mathbb{R}^d}\mathrm{tr}(\Omega Q^2 - Q\Omega Q)\,dx}_{\overset{\mathrm{def}}{=}\mathcal{I}}$$

$$\underbrace{+\xi\int_{\mathbb{R}^d} D_{\alpha\gamma}(Q_{\gamma\beta}+\frac{\delta_{\gamma\beta}}{d})Q_{\alpha\beta}+(Q_{\alpha\gamma}+\frac{\delta_{\alpha\gamma}}{d})D_{\gamma\beta}Q_{\alpha\beta}-2(Q_{\alpha\beta}+\frac{\delta_{\alpha\beta}}{d})Q_{\alpha\beta}\mathrm{tr}(Q\nabla u)\,dx}_{\overset{\text{def}}{=}\mathcal{II}}$$

Recalling that Q is symmetric we have $\mathcal{I}=0$. Also:

$$|\mathcal{II}|=|2\xi||\int_{\mathbb{R}^d}\frac{1}{d}D_{\alpha\beta}Q_{\alpha\beta}+D_{\alpha\gamma}Q_{\gamma\beta}Q_{\beta\alpha}-Q_{\alpha\beta}Q_{\alpha\beta}\mathrm{tr}(Q\nabla u)\,dx|\le C(d)\int_{\mathbb{R}^d}\varepsilon|\nabla u|^2+\int_{\mathbb{R}^d}\frac{|\xi|^2}{\varepsilon}(|Q|^2+|Q|^6)\,dx$$

Thus we get:

$$\frac{d}{dt}\int_{\mathbb{R}^d}|Q|^2\,dx\le C(d)\varepsilon\int_{\mathbb{R}^d}|\nabla u|^2\,dx+\frac{|\xi|^2}{\varepsilon}\int_{\mathbb{R}^d}|Q|^2+|Q|^6\,dx+\hat{C}\int_{\mathbb{R}^d}|Q|^2+|Q|^4\,dx \tag{23}$$

with $\hat{C}=\hat{C}(a,b)>0$.

Let us observe now that there exists $M=M(a,b,c)$ large enough, so that

$$\frac{M}{2}\mathrm{tr}(Q^2)+\frac{c}{8}\mathrm{tr}^2(Q^2)\le(M+\frac{a}{2})\mathrm{tr}(Q^2)-\frac{b}{3}\mathrm{tr}(Q^3)+\frac{c}{4}\mathrm{tr}^2(Q^2) \tag{24}$$

for any $Q\in S_0$.

Multiplying the equation (23) by M and adding to (22) we obtain:

$$\begin{aligned}&\frac{d}{dt}(E(t)+M\|Q\|^2_{L^2_x})+\nu\|\nabla u\|^2_{L^2_x}+\Gamma\lambda L^2\|\Delta Q\|^2_{L^2_x}+\Gamma\lambda c^2\|Q\|^6_{L^6}+a^2\|Q\|^2_{L^2_x}\\&\qquad+2cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla Q|^2\mathrm{tr}(Q^2)\,dx+cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla\left(\mathrm{tr}(Q^2)\right)|^2\,dx\\&\le\bar{C}\left(\|\nabla Q\|^2_{L^2_x}+\|Q\|^4_{L^4}\right)+MC(d)\varepsilon\int_{\mathbb{R}^d}|\nabla u|^2\,dx+\frac{M|\xi|^2}{\varepsilon}\int_{\mathbb{R}^d}|Q|^2+|Q|^6\,dx+M\hat{C}\int_{\mathbb{R}^d}|Q|^2+|Q|^4\,dx\end{aligned} \tag{25}$$

We chose ε small enough so that $MC(d)\varepsilon<\nu$. Finally we make the assumption that $|\xi|$ is small enough, depending on a,b,c,d,ν so that

$$\frac{M|\xi|^2}{\varepsilon}\le\Gamma\lambda c^2$$

Then taking into account equation (24) we obtain the claimed relation (16).□

We note that the ξ small hypothesis is necessary because we are in infinite domain, for example, in the periodic domain, we can add a constant to the functional and get the apriori L^p estimates without any smallness condition on ξ.

3. The existence of weak solutions

The next proposition follows closely the similar result in [36] where it was done for $\lambda=1$. The purpose for including it here is to provide an alternative approximation system thus correcting the proof in [36] and also to show how the cancellations that appeared previously in the derivation of the energy law still survive at the approximate level but with some differences, phenomenon which will appear in a much more complex setting in the proof of uniqueness in the next section.

Proposition 3.1. *For $d=2,3$ there exists a weak solution (Q,u) of the system* (8) *subject to initial conditions* (9)*. The solution (Q,u) is such that $Q\in L^\infty_{loc}(\mathbb{R}_+;H^1)\cap L^2_{loc}(\mathbb{R}_+;H^2)$ and $u\in L^\infty_{loc}(\mathbb{R}_+;L^2)\cap L^2_{loc}(\mathbb{R}_+;H^1)$.*

Proof. As first step of the construction of weak solutions for the system (8) we construct for any fixed $\varepsilon>0$ a global weak solution

$$Q_\varepsilon\in L^\infty_{loc}(\mathbb{R}_+;H^1)\cap L^2_{loc}(\mathbb{R}_+;H^2)\quad u_\varepsilon\in L^\infty_{loc}(\mathbb{R}_+;L^2)\cap L^2_{loc}(\mathbb{R}_+;H^1)$$

for the modified system obtained by mollifying the coefficients of the equation for the Q tensor and by adding to the equation of the velocity a regularizing term. This term is needed in order to estimate some "bad" terms which does not disappear in an energy estimate. For the simplicity of the notations, we drop the indices ε

and we denote the solution $(Q_\varepsilon, u_\varepsilon)$ by (Q, u).

$$\begin{cases}
\partial_t Q + (R_\varepsilon u)\nabla Q - \Big(\big(R_\varepsilon(\xi D + \Omega)\big)(Q + \frac{1}{d}Id)\Big) \\
-\Big((Q + \frac{1}{d}Id)R_\varepsilon(\xi D - \Omega)\Big) \\
+2\xi\Big((Q + \frac{1}{d}Id)\mathrm{tr}\big(Q\nabla R_\varepsilon u\big)\Big) = \Gamma H \\
\partial_t u + (R_\varepsilon u)\nabla u - \nu\Delta u + \nabla p = -\varepsilon\mathcal{P}R_\varepsilon\left(\sum_{l,m=1}^d \nabla Q_{lm}\left(R_\varepsilon u\cdot\nabla Q_{lm}\right)|R_\varepsilon u\nabla Q|\right) \\
+\varepsilon\mathcal{P}\nabla\cdot R_\varepsilon\Bigg(\nabla R_\varepsilon u|\nabla R_\varepsilon u|^2\Bigg) - \lambda\xi\nabla\cdot R_\varepsilon\Bigg(\left(Q + \frac{1}{d}Id\right)H\Bigg) - \xi\mathcal{P}\nabla\cdot R_\varepsilon\Bigg(H\left(Q + \frac{1}{d}Id\right)\Bigg) \\
+2\lambda\xi\nabla\cdot R_\varepsilon\Bigg(\big(Q + \frac{1}{d}Id\big)\big(QH\big)\Bigg) - L\lambda R_\varepsilon(\nabla\cdot\mathrm{tr}(\nabla Q\nabla Q)) \\
+L\lambda\mathcal{P}\nabla\cdot R_\varepsilon\left(Q\Delta Q - \Delta QQ\right) \\
(Q, u)|_{t=0} = (R_\varepsilon\overline{Q}, R_\varepsilon\overline{u}).
\end{cases}$$

where R_ε is the convolution operator with the kernel $\epsilon^{-d}\chi(\epsilon^{-1}\cdot)$.

In order to construct the global weak solution for this system, we use the classical Friedrich's scheme. We define the mollifying operator

$$\widehat{J_n f}(\xi) \overset{\text{def}}{=} 1_{\{2^{-n}\leq|\xi|\leq 2^n\}}\hat{f}(\xi).$$

We consider the approximating system:

$$\begin{cases}
\partial_t Q^{(n)} + J_n\Big(R_\varepsilon J_n u^n\nabla J_n Q^{(n)}\Big) - J_n\Big((\xi J_n R_\varepsilon D^{(n)} + J_n R_\varepsilon\Omega^{(n)})(J_n Q^{(n)} + \frac{1}{d}Id)\Big) \\
-J_n\Big((J_n Q^{(n)} + \frac{1}{d}Id)(\xi J_n R_\varepsilon D^{(n)} - J_n R_\varepsilon\Omega^{(n)})\Big) \\
+2\xi J_n\Big((J_n Q^{(n)} + \frac{1}{d}Id)\mathrm{tr}\big(J_n Q^{(n)}\nabla J_n R_\varepsilon u^{(n)}\big)\Big) = \Gamma\tilde{H}^{(n)} \\
\partial_t u^n + \mathcal{P}J_n(\mathcal{P}J_n R_\varepsilon u^n\nabla\mathcal{P}J_n u^n) - \nu\Delta\mathcal{P}J_n u^{(n)} = \\
-\varepsilon\mathcal{P}J_n R_\varepsilon\Bigg(\sum_{l,m=1}^d \nabla J_n Q^{(n)}_{lm}\left(R_\varepsilon J_n u^n\cdot\nabla J_n Q^{(n)}_{lm}\right)|R_\varepsilon J_n u^n\nabla J_n Q^{(n)}|\Bigg) \\
+\varepsilon\mathcal{P}\nabla\cdot J_n R_\varepsilon\Bigg(\nabla R_\varepsilon J_n u^{(n)}|\nabla R_\varepsilon J_n u^{(n)}|^2\Bigg) \\
-\lambda\xi\mathcal{P}\nabla\cdot J_n\Bigg(\left(J_n Q^{(n)} + \frac{1}{d}Id\right)\tilde{H}^{(n)}\Bigg) - \lambda\xi\mathcal{P}\nabla\cdot J_n\Bigg(\tilde{H}^{(n)}\left(J_n Q^{(n)} + \frac{1}{d}Id\right)\Bigg) \\
+2\lambda\xi\mathcal{P}\nabla\cdot J_n\Bigg(\big(J_n Q^{(n)} + \frac{1}{d}Id\big)\big(J_n Q^{(n)}\tilde{H}^{(n)}\big)\Bigg) - L\lambda\mathcal{P}J_n(\nabla\cdot\mathrm{tr}(J_n Q^{(n)}\nabla J_n Q^{(n)})) \\
+L\lambda\mathcal{P}\nabla\cdot J_n\left(J_n Q^{(n)}\Delta J_n Q^{(n)} - \Delta J_n Q^{(n)} J_n Q^{(n)}\right)
\end{cases} \tag{26}$$

where $\mathcal{P}$ denotes the Leray projector onto divergence-free vector fields, M is a positive constant, and $\tilde{H}^{(n)} \overset{def}{=} L\Delta J_n Q^{(n)} - aJ_n Q^{(n)} + bJ_n[(J_n Q^{(n)} J_n Q^{(n)}) - \frac{\mathrm{tr}(J_n Q^{(n)} J_n Q^{(n)})}{d}Id] - cJ_n\Bigg(J_n Q^{(n)}\big|J_n Q^{(n)}\big|^2\Bigg)$. We take as initial data $(J_n R_\varepsilon\bar{Q}, J_n R_\varepsilon\bar{u})$.

The system above can be regarded as an ordinary differential equation in L^2 verifying the conditions of the Cauchy-Lipschitz theorem. Thus it admits a unique maximal solution $(Q^{(n)}, u^{(n)}) \in C^1([0, T_n); L^2(\mathbb{R}^d; \mathbb{R}^{d\times d})\times L^2(\mathbb{R}^d, \mathbb{R}^d))$. As we have $(\mathcal{P}J_n)^2 = \mathcal{P}J_n$ and $J_n^2 = J_n$ the pair $(J_n Q^{(n)}, \mathcal{P}J_n u^{(n)})$ is also a solution of (26). By uniqueness we have $(J_n Q^{(n)}, \mathcal{P}J_n u^{(n)}) = (Q^{(n)}, u^{(n)})$ hence $(Q^{(n)}, u^{(n)}) \in C^1([0, T_n), H^\infty)$ and $(Q^{(n)}, u^{(n)})$ satisfy the system:

$$\begin{cases}
\partial_t Q^{(n)} + J_n\Big(R_\varepsilon u^n\nabla Q^{(n)}\Big) - J_n\Big((\xi R_\varepsilon D^{(n)} + R_\varepsilon\Omega^{(n)})(Q^{(n)} + \frac{1}{d}Id)\Big) \\
-J_n\Big((Q^{(n)} + \frac{1}{d}Id)(\xi R_\varepsilon D^{(n)} - R_\varepsilon\Omega^{(n)})\Big) + 2\xi J_n\Big((Q^{(n)} + \frac{1}{d}Id)\mathrm{tr}\big(Q^{(n)}\nabla R_\varepsilon u^n\big)\Big) = \Gamma\bar{H}^{(n)} \\
\partial_t u^n + \mathcal{P}J_n(R_\varepsilon u^n\nabla u^n) - \nu\Delta u^{(n)} = \\
-\varepsilon\mathcal{P}J_n\Bigg(\sum_{l,m=1}^d \nabla Q^{(n)}_{lm}\left(R_\varepsilon u^n\cdot\nabla Q^{(n)}_{lm}\right)|R_\varepsilon u^n\nabla Q^{(n)}|\Bigg) \\
+\varepsilon\mathcal{P}\nabla\cdot J_n R_\varepsilon\Bigg(\nabla R_\varepsilon u^{(n)}|\nabla R_\varepsilon u^{(n)}|^2\Bigg) \\
-\lambda\xi\mathcal{P}\nabla\cdot J_n\Bigg(\left(Q^{(n)} + \frac{1}{d}Id\right)\bar{H}^{(n)}\Bigg) - \lambda\xi\mathcal{P}\nabla\cdot J_n\Bigg(\bar{H}^{(n)}\left(Q^{(n)} + \frac{1}{d}Id\right)\Bigg) \\
+2\lambda\xi\mathcal{P}\nabla\cdot J_n\Bigg(\big(Q^{(n)} + \frac{1}{d}Id\big)\big(Q^{(n)}\bar{H}^{(n)}\big)\Bigg) \\
-L\lambda\mathcal{P}J_n(\nabla\cdot\mathrm{tr}(\nabla Q^{(n)}\nabla Q^{(n)})) + L\lambda\mathcal{P}\nabla\cdot J_n\left(Q^{(n)}\Delta Q^{(n)} - \Delta Q^{(n)}Q^{(n)}\right)
\end{cases} \tag{27}$$

where $\bar{H}^{(n)} \overset{def}{=} L\Delta Q^{(n)} - aQ^{(n)} + bJ_n[(Q^{(n)}Q^{(n)}) - \frac{\mathrm{tr}(J_n(Q^{(n)}Q^{(n)}))}{d} Id] - cJ_n(Q^{(n)}|Q^{(n)}|^2)$. The initial data is $(J_n\bar{Q}, J_n\bar{u})$. We recall now a few properties of J_n :

Lemma 3.2. *The operators $\mathcal{P}$ and J_n are selfadjoint in L^2. Moreover J_n and $\mathcal{P}J_n$ are also idempotent and J_n commutes with distributional derivatives.*

We proceed in a manner analogous to the proof of Proposition 2.1 and multiply the first equation in (27) by $-\lambda\bar{H}^{(n)}$, take the trace, integrate over $\mathbb{R}^d$ and by parts, and add to the second equation multiplied by $u^{(n)}$. Let us observe that almost all the cancellations in the proof of (2.1) hold, except for a few terms that need to be estimated separately. We also have some more new terms that we added in the regularization, terms that control the ones which do not cancel. Thus we have:

$$
\begin{aligned}
&\frac{d}{dt}\int_{\mathbb{R}^d} \frac{1}{2}|u^n|^2 + \frac{L\lambda}{2}|\nabla Q^{(n)}|^2 + \lambda\big(\frac{a}{2}|Q^{(n)}|^2 - \frac{b}{3}\mathrm{tr}(Q^{(n)})^3 + \frac{c}{4}|Q^{(n)}|^4\big)\,dx\\
&+\nu\int_{\mathbb{R}^d}|\nabla u^n|^2\,dx + \Gamma\lambda\int_{\mathbb{R}^d}\mathrm{tr}\Big[J_n\Big(L\Delta Q^{(n)} - aQ^{(n)} + b[(Q^{(n)})^2 - \frac{\mathrm{tr}((Q^{(n)})^2)}{3}Id] - cQ^{(n)}\big|Q^{(n)}\big|^2\Big)\Big]^2\,dx\\
&+\varepsilon\int_{\mathbb{R}^d}|R_\varepsilon u\nabla Q^{(n)}|^3\,dx + \varepsilon\int_{\mathbb{R}^d}|R_\varepsilon\nabla u^n|^4\,dx\\
&\leq \lambda\int_{\mathbb{R}^d} J_n\left(R_\varepsilon u^n\cdot\nabla Q^{(n)}_{\alpha\beta}\right) J_n\left(bQ^{(n)}_{\alpha\gamma}Q^{(n)}_{\gamma\beta} - cQ^{(n)}_{\alpha\beta}\big|Q^{(n)}\big|^2\right)\,dx\\
&+\lambda\int_{\mathbb{R}^d} J_n\left(-R_\varepsilon\Omega^{(n)}_{\alpha\gamma}Q^{(n)}_{\gamma\beta} + Q^{(n)}_{\alpha\gamma}R_\varepsilon\Omega^{(n)}_{\gamma\beta}\right) J_n\left(bQ^{(n)}_{\alpha\delta}Q^{(n)}_{\delta\beta} - cQ^{(n)}_{\alpha\beta}\big|Q^{(n)}\big|^2\right)\,dx \qquad (28)
\end{aligned}
$$

hence :

$$
\begin{aligned}
&\frac{d}{dt}\int_{\mathbb{R}^d} \frac{1}{2}|u^n|^2 + \frac{L\lambda}{2}|\nabla Q^{(n)}|^2 + \lambda\big(\frac{a}{2}|Q^{(n)}|^2 - \frac{b}{3}\mathrm{tr}(Q^{(n)})^3 + \frac{c}{4}|Q^{(n)}|^4\big)\,dx + \nu\int_{\mathbb{R}^d}|\nabla u^n|^2\,dx\\
&+\Gamma\lambda\int_{\mathbb{R}^d} L^2|\Delta Q^{(n)}|^2\,dx + \Gamma\lambda a^2\int_{\mathbb{R}^d}|Q^{(n)}|^2\,dx + C(b^2,d,\Gamma,\lambda)\int_{\mathbb{R}^d}|Q^{(n)}|^4\,dx + \Gamma\lambda c^2\int_{\mathbb{R}^d}\big|J_n(Q^{(n)}|Q^{(n)}|^2)\big|^2\,dx\\
&+\varepsilon\int_{\mathbb{R}^d}|R_\varepsilon u\nabla Q^{(n)}|^3\,dx + \varepsilon\int_{\mathbb{R}^d}|R_\varepsilon\nabla u^n|^4\,dx\\
&\leq \underbrace{2\Gamma\lambda c\int_{\mathbb{R}^d} L\Delta Q^{(n)}\cdot Q^{(n)}|Q^{(n)}|^2\,dx}_{\overset{\mathrm{def}}{=}\mathcal{I}} - 2\Gamma\lambda\int_{\mathbb{R}^d} L\Delta Q^{(n)}\cdot\big(-aQ^{(n)} + bJ_n\Big[(Q^{(n)})^2 - \frac{\mathrm{tr}(Q^{(n)})^2}{d}Id)\Big]\big)\,dx\\
&-2\Gamma\lambda\int_{\mathbb{R}^d} cQ^{(n)}|Q^{(n)}|^2\cdot\left(aQ^{(n)} - bJ_n\Big[(Q^{(n)})^2 - \frac{\mathrm{tr}(Q^{(n)})^2}{d}Id\Big]\right)\,dx\\
&+\lambda\underbrace{\int_{\mathbb{R}^d} J_n\left(R_\varepsilon u^n\cdot\nabla Q^{(n)}_{\alpha\beta}\right) J_n\left(bQ^{(n)}_{\alpha\gamma}Q^{(n)}_{\gamma\beta} - cQ^{(n)}_{\alpha\beta}\big|Q^{(n)}\big|^2\right)\,dx}_{\overset{\mathrm{def}}{=}\mathcal{II}}\\
&+C\int_{\mathbb{R}^d}|R_\varepsilon\nabla u^n|^2|Q^{(n)}|^2\,dx + \frac{\Gamma c^2}{8}\int_{\mathbb{R}^d}|J_n(Q^{(n)}|Q^{(n)}|^2)|^2\,dx + C\int_{\mathbb{R}^d}|Q^{(n)}|^4\,dx
\end{aligned}
$$

We have that

$$
\begin{aligned}
\mathcal{II} &= \int_{\mathbb{R}^d}\left(R_\varepsilon u^n\cdot\nabla Q^{(n)}_{\alpha\beta}\right) J_n\left(bQ^{(n)}_{\alpha\gamma}Q^{(n)}_{\gamma\beta} - cQ^{(n)}_{\alpha\beta}\big|Q^{(n)}\big|^2\right)\,dx\\
&\leq \left(\frac{4}{\Gamma c^2} + \frac{1}{4C(b^2,d,\Gamma)}\right)\int_{\mathbb{R}^d}|R_\varepsilon u^n\cdot\nabla Q^{(n)}|^2\,dx\\
&+ \frac{C(b^2,d,\Gamma)}{2}\|Q^{(n)}\|^4_{L^4} + \frac{\Gamma c^2}{8}\int_{\mathbb{R}^d}|J_n(Q^{(n)}|Q^{(n)}|^2)|^2\,dx\\
&\leq \frac{\varepsilon}{2}\int_{\mathbb{R}^d}|R_\varepsilon u^n\cdot\nabla Q^{(n)}|^3 + C(\varepsilon,b^2,c^2,d,\Gamma)\int_{\mathbb{R}^d}\sum_{l,m=1}^{d}|R_\varepsilon u^n\cdot\nabla Q^{(n)}_{lm}|\,dx \qquad (29)\\
&+ \frac{C(b^2,c^2,d,\Gamma)}{2}\|Q^{(n)}\|^4_{L^4} + \frac{\Gamma c^2}{8}\int_{\mathbb{R}^d}|J_n(Q^{(n)}|Q^{(n)}|^2)|^2\,dx\\
&\leq \frac{\varepsilon}{2}\int_{\mathbb{R}^d}|R_\varepsilon u^n\cdot\nabla Q^{(n)}|^3\,dx + C_1(\varepsilon,b^2,c^2,d,\Gamma)\int_{\mathbb{R}^d}|u^n|^2\,dx + C_2(\varepsilon,b,c,d^2,\Gamma)\int_{\mathbb{R}^d}|\nabla Q^{(n)}|^2\,dx\\
&+ \frac{C(b^2,d,\Gamma)}{2}\|Q^{(n)}\|^4_{L^4} + \frac{\Gamma c^2}{8}\int_{\mathbb{R}^d}|J_n(Q^{(n)}|Q^{(n)}|^2)|^2\,dx
\end{aligned}
$$

Using the fact that $\mathcal{I} \leq 0$ and the estimate for $\mathcal{II}$ shown before, we replace in (28) and obtain:

$$\begin{aligned} &\frac{d}{dt}\int_{\mathbb{R}^d} \frac{1}{2}|u^n|^2 + \frac{L\lambda}{2}|\nabla Q^{(n)}|^2 + \lambda\big(\frac{a}{2}|Q^{(n)}|^2 - \frac{b}{3}\mathrm{tr}(Q^{(n)})^3 + \frac{c}{4}|Q^{(n)}|^4\big)\,dx \\ +\nu\int_{\mathbb{R}^d}|\nabla u^n|^2\,dx + \Gamma\lambda\int_{\mathbb{R}^d} L^2|\Delta Q^{(n)}|^2\,dx + \frac{\varepsilon}{2}\int_{\mathbb{R}^d}|R_\varepsilon u^n\cdot\nabla Q^{(n)}|^3\,dx + \frac{\varepsilon}{2}\int_{\mathbb{R}^d}|\nabla R_\varepsilon u^n|^4\,dx \\ &\leq \int_{\mathbb{R}^d}|Q^{(n)}|^2 + |Q^{(n)}|^4\,dx + C\int_{\mathbb{R}^d}|\nabla Q^{(n)}|^2\,dx + C(\varepsilon)\int_{\mathbb{R}^d}|u^{(n)}|^2\,dx \end{aligned}$$

This estimate does not readily provide bounds on $Q^{(n)}$ because the term $\frac{a}{2}|Q^{(n)}|^2 - \frac{b}{3}\mathrm{tr}(Q^{(n)})^3 + \frac{c}{4}|Q^{(n)}|^4$ could be negative. In order to obtain H^1 estimates we proceed as in the proof of Proposition 2.3. We put the proof in the appendix by Proposition B.1. We can continue to proceed as in the proof of Proposition 2.3 and in fact in this case because of the first two regularizing terms on the right hand side of the u^n equation in (27) we do not need the ξ small assumption. These estimates allow us to conclude that $T_n = \infty$ and we also get the following apriori bounds:

$$\begin{aligned} \sup_n \|\nabla R_\varepsilon u^n\|_{L^4(0,T;L^4)}, \sup_n \|R_\varepsilon u^n\cdot\nabla Q^{(n)}\|_{L^3(0,T;L^3)}, dx \leq C(\varepsilon) \\ \sup_n \|Q^{(n)}\|_{L^2(0,T;H^2)\cap L^\infty(0,T;H^1)} < \infty, \\ \sup_n \|u^n\|_{L^\infty(0,T;L^2)\cap L^2(0,T;H^1)} < \infty, \end{aligned} \tag{30}$$

for any $T < \infty$. By the bounds which can be obtained by using the equation on $\partial_t(Q^{(n)}, u^n)$ in some $L^\infty_{loc}(H^{-N})$ for large enough N, we get, by classical local compactness Aubin-Lions lemma, on a subsequence, that:

$$\begin{aligned} Q^{(n)} \rightharpoonup Q \text{ in } L^2(0,T;H^2) \text{ and } Q^{(n)} \to Q \text{ in } L^2(0,T;H^{2-\delta}_{loc}), \forall\delta > 0 \\ Q^{(n)}(t) \rightharpoonup Q(t) \text{ in } H^1 \text{ for all } t\in\mathbb{R}_+ \\ u^n \rightharpoonup u \text{ in } L^2(0,T;H^1) \text{ and } u^n \to u \text{ in } L^2(0,T;H^{1-\delta}_{loc}), \forall\delta > 0 \\ u^n(t) \rightharpoonup u(t) \text{ in } L^2 \text{ for all } t\in\mathbb{R}_+ \end{aligned}$$

Thus we can pass to the limit and obtain a weak solution of the approximating system:

$$\begin{cases} \partial_t Q^{(\varepsilon)} + R_\varepsilon u^\varepsilon\nabla Q^{(\varepsilon)} - (\xi R_\varepsilon D^\varepsilon + R_\varepsilon\Omega^\varepsilon)(Q^{(\varepsilon)} + \frac{1}{d}Id)\Big) + \Big((Q^{(\varepsilon)} + \frac{1}{d}Id)(\xi R_\varepsilon D^\varepsilon - R_\varepsilon\Omega^\varepsilon)\Big) \\ -2\xi\Big((Q^{(\varepsilon)} + \frac{1}{d}Id)\mathrm{tr}\big(Q^{(\varepsilon)}\nabla u^\varepsilon\big)\Big) = \Gamma H^\varepsilon \\ \partial_t u^\varepsilon + \mathcal{P}R_\varepsilon u^\varepsilon\nabla u^\varepsilon = -\varepsilon\mathcal{P}R_\varepsilon\left(\sum_{l,m=1}^d \nabla Q_{lm}\left(R_\varepsilon u\cdot\nabla Q_{lm}\right)|R_\varepsilon u\nabla Q|\right) \\ +\varepsilon\mathcal{P}\nabla\cdot R_\varepsilon\Big(R_\varepsilon\nabla u|R_\varepsilon\nabla u|^2\Big) - \lambda\xi\mathcal{P}\nabla\cdot R_\varepsilon\Big(\left(Q^{(\varepsilon)} + \frac{1}{d}Id\right)H^\varepsilon\Big) \\ -\lambda\xi\mathcal{P}\nabla\cdot R_\varepsilon\Big(H^\varepsilon\left(Q^{(\varepsilon)} + \frac{1}{d}Id\right)\Big) + 2\lambda\xi\mathcal{P}\nabla\cdot R_\varepsilon\Big(\big(Q^{(\varepsilon)} + \frac{1}{d}\big)\big(Q^{(\varepsilon)}H^\varepsilon\big)\Big) \\ -L\lambda\mathcal{P}(\nabla\cdot R_\varepsilon\mathrm{tr}(\nabla Q^{(\varepsilon)}\odot\nabla Q^{(\varepsilon)})) + L\lambda\mathcal{P}\nabla\cdot R_\varepsilon\left(Q^{(\varepsilon)}\Delta Q^{(\varepsilon)} - \Delta Q^{(\varepsilon)}Q^{(\varepsilon)}\right) + \nu\Delta u^\varepsilon \end{cases} \tag{31}$$

where we recall that $H = L\Delta Q^{(\varepsilon)} - aQ^{(\varepsilon)} + b[(Q^{(\varepsilon)})^2 - \frac{\mathrm{tr}((Q^{(\varepsilon)})^2))}{d}Id] - cQ^{(\varepsilon)}\mathrm{tr}((Q^{(\varepsilon)})^2)$. The initial data for the limit system is $(R_\varepsilon\bar{Q}, R_\varepsilon\bar{u})$.

One can easily see that the solutions of (31) are smooth, first by obtaining C^∞ regularity for the first Q equations, by bootstrapping the regularity improvement provided by the linear heat equation, and then the regularity for the u equation, by bootstrapping the regularity improvement provided by a linear advection equation. For this system we can proceed as in the case of apriori estimates and obtain the same estimates, independent of ε because the solutions are smooth and all the cancellations that were used in the apriori estimates also hold here. In particular we obtain:

$$\begin{aligned} \sup_\varepsilon \|Q^{(\varepsilon)}\|_{L^\infty(0,T;H^1)\cap L^2(0,T;H^2)} < \infty, \\ \sup_\varepsilon \|u^\varepsilon\|_{L^\infty(0,T;L^2)\cap L^2(0,T;H^1)} < \infty \end{aligned} \tag{32}$$

for any $T < \infty$. Taking into account those bounds and also the bounds which can be obtained by using the equation on $\partial_t(Q^\varepsilon, u^\varepsilon)$ in some $L^p_{loc}(H^{-N})$ for large enough N, we get, by classical local compactness Aubin-Lions lemma and by weak convergence arguments, that there exists a $Q\in L^\infty_{loc}(\mathbb{R}_+;H^1)\cap L^2_{loc}(\mathbb{R}_+;H^2)$ and a $u\in L^\infty_{loc}(\mathbb{R}_+;L^2)\cap L^2_{loc}(\mathbb{R}_+;H^1)$ so that, on a subsequence, we have:

$$Q^{(\varepsilon)} \rightharpoonup Q \text{ in } L^2(0,T;H^2) \text{ and } Q^{(n)} \to Q \text{ in } L^2(0,T;H^{2-\delta}_{loc}), \forall\delta > 0$$

$$
\begin{aligned}
Q^{(\varepsilon)}(t) \rightharpoonup Q(t) \text{ in } H^1 \text{ for all } t\in\mathbb{R}_+\\
u^\varepsilon \rightharpoonup u \text{ in } L^2(0,T;H^1) \text{ and } u^n \to u \text{ in } L^2(0,T;H^{1-\delta}_{loc}), \forall \delta>0\\
u^\varepsilon(t) \rightharpoonup u(t) \text{ in } L^2 \text{ for all } t\in\mathbb{R}_+
\end{aligned}
\tag{33}
$$

These convergences allow us to the pass to the limit in the weak solutions of the system (31) to obtain a weak solution of (8), namely (10),(11). Of all the terms there is only one type that is slightly difficult to treat in passing to the limit, namely:

$$
L\int_0^\infty\int_{\mathbb{R}^d}\partial_\beta\left(Q^{(\varepsilon)}_{\alpha\gamma}\Delta Q^{(\varepsilon)}_{\gamma\beta}-\Delta Q^{(\varepsilon)}_{\alpha\gamma}Q^{(\varepsilon)}_{\gamma\beta}\right)\psi_\alpha\,dx\,dt=-L\int_0^\infty\int_{\mathbb{R}^d}\left(Q^{(\varepsilon)}_{\alpha\gamma}\Delta Q^{(\varepsilon)}_{\gamma\beta}-\Delta Q^{(\varepsilon)}_{\alpha\gamma}Q^{(\varepsilon)}_{\gamma\beta}\right)\cdot\psi_{\alpha,\beta}\,dx\,dt.
$$

Taking into account that ψ is compactly supported and the convergences (33) one can easily pass to the limit the terms $\psi_{\alpha,\beta}Q^{(\varepsilon)}_{\alpha\gamma}$ and $\psi_{\alpha,\beta}Q^{(\varepsilon)}_{\gamma\beta}$ strongly in $L^2(0,T;L^2)$. Relations (33) give that $\Delta Q^{(\varepsilon)}_{\gamma\beta}$, $\Delta Q^{(\varepsilon)}_{\alpha\gamma}$ converges weakly in $L^2(0,T;L^2)$. Thus we get convergence to the limit term

$$
\begin{aligned}
&L\int_0^\infty\int_{\mathbb{R}^d}\partial_\beta(Q_{\alpha\gamma}\Delta Q_{\gamma\beta})\psi_\alpha dxdt-L\int_0^\infty\int_{\mathbb{R}^d}\partial_\beta(\Delta Q_{\alpha\gamma})Q_{\gamma\beta})\psi_\alpha dxdt\\
&=-L\int_0^T\int_{\mathbb{R}^d}(\Delta Q_{\gamma\beta})(\partial_\beta\psi_\alpha Q_{\alpha\gamma})dxdt+L\int_0^T\int_{\mathbb{R}^d}(\Delta Q_{\alpha\gamma})(\partial_\beta\psi_\alpha Q_{\gamma\beta})dxdt.
\end{aligned}
\tag{34}
$$

Using also the uniform bound of $\varepsilon\|R_\varepsilon u^\varepsilon\nabla Q^\varepsilon\|^3_{L^3}$ it is easy to check that $\varepsilon\int|R_\varepsilon u^\varepsilon\nabla Q^\varepsilon|^2\nabla Q^\varepsilon\cdot R_\varepsilon\mathcal{P}\varphi dxdt$ converges to zero. A similar observation holds for the ε-regularisation term $\varepsilon\mathcal{P}\nabla\cdot\Bigg(R_\varepsilon\nabla u|R_\varepsilon\nabla u|^2\Bigg)$. $\square$

4. The uniqueness of weak solutions

We start with a number of technical tools that are crucial for our proof.

4.1. Littlewood-Paley theory. We define $\mathcal{C}$ to be the ring of center 0, of small radius 1/2 and great radius 2. There exist two nonnegative radial functions χ and φ belonging respectively to $\mathcal{D}(B(0,1))$ and to $\mathcal{D}(\mathcal{C})$ so that

$$
\chi(\xi)+\sum_{q\geq 0}\varphi(2^{-q}\xi)=1,\forall\xi\in\mathbb{R}^d \tag{35}
$$

$$
|p-q|\geq 5\Rightarrow \mathrm{Supp}\ \varphi(2^{-q}\cdot)\cap\mathrm{Supp}\ \varphi(2^{-p}\cdot)=\emptyset. \tag{36}
$$

For instance, one can take $\chi\in\mathcal{D}(B(0,1))$ such that $\chi\equiv 1$ on $B(0,1/2)$ and take

$$
\varphi(\xi)=\chi(\xi/2)-\chi(\xi).
$$

Then, we are able to define the Littlewood-Paley decomposition. Let us denote by $\mathcal{F}$ the Fourier transform on $\mathbb{R}^d$. Let $h,\ \tilde{h},\ \dot{\Delta}_q, \dot{S}_q$ $(q\in\mathbb{Z})$ be defined as follows:

$$
\begin{aligned}
h=\mathcal{F}^{-1}\varphi \quad &\text{and} \quad \tilde{h}=\mathcal{F}^{-1}\chi,\\
\dot{\Delta}_q u=\mathcal{F}^{-1}(\varphi(2^{-q}\xi)\mathcal{F}u)&=2^{qd}\int h(2^q y)u(x-y)dy,\\
\dot{S}_q u=\mathcal{F}^{-1}(\chi(2^{-q}\xi)\mathcal{F}u)&=2^{qd}\int \tilde{h}(2^q y)u(x-y)dy.
\end{aligned}
$$

We recall that for two appropriately smooth functions a and b we have the Bony's paraproduct decomposition [4]:

$$
ab=\dot{T}_a b+\dot{T}_b a+\dot{R}(a,b)
$$

where

$$
\dot{T}_a b=\sum_{q\in\mathbb{Z}}\dot{S}_{q-1}a\dot{\Delta}_q b,\quad \dot{T}_b a=\sum_{q\in\mathbb{Z}}\dot{S}_{q-1}b\dot{\Delta}_q a,\quad\text{and}\quad \dot{R}(a,b)=\sum_{\substack{q\in\mathbb{Z},\\ i\in\{0,\pm1\}}}\dot{\Delta}_q a\dot{\Delta}_{q+i}b.
$$

Then we have

$$
\dot{\Delta}_q(ab)=\dot{\Delta}_q\dot{T}_a b+\dot{\Delta}_q\dot{T}_b a+\dot{\Delta}_q\dot{R}(a,b)=\dot{\Delta}_q\dot{T}_a b+\dot{\Delta}_q\tilde{R}(a,b),
$$

where $\tilde{R}(a,b) = \dot{T}_b a + \dot{R}(a,b) = \sum_{q\in\mathbb{Z}} \dot{S}_{q+2} b \dot{\Delta}_q a$. Moreover:

$$\begin{aligned}
\dot{\Delta}_q(ab) &= \sum_{|q'-q|\leq 5} \dot{\Delta}_q(\dot{S}_{q'-1}a\dot{\Delta}_{q'}b) + \sum_{q'>q-5} \dot{\Delta}_q(\dot{S}_{q'+2}b\dot{\Delta}_{q'}a)\\
&= \sum_{|q'-q|\leq 5} [\dot{\Delta}_q,\, \dot{S}_{q'-1}a]\dot{\Delta}_{q'}b + \sum_{|q'-q|\leq 5} \dot{S}_{q'-1}a\dot{\Delta}_q\dot{\Delta}_{q'}b + \sum_{q'>q-5} \dot{\Delta}_q(S_{q'+2}b\dot{\Delta}_{q'}a)\\
&= \sum_{|q'-q|\leq 5} [\dot{\Delta}_q,\, \dot{S}_{q'-1}a]\dot{\Delta}_{q'}b + \sum_{|q'-q|\leq 5} (\dot{S}_{q'-1}a - \dot{S}_{q-1}a)\dot{\Delta}_q\dot{\Delta}_{q'}b\\
&\qquad\qquad + \sum_{q'>q-5} \dot{\Delta}_q(\dot{S}_{q'+2}b\dot{\Delta}_{q'}a) + \underbrace{\sum_{|q'-q|\leq 5} \dot{S}_{q-1}a\dot{\Delta}_q\dot{\Delta}_{q'}b}_{=\dot{S}_{q-1}a\dot{\Delta}_q b}
\end{aligned} \tag{37}$$

In terms of this decomposition we can express the Sobolev norm of an element u in the (nonhomogeneous!) space H^s as:

$$\|u\|_{H^s} = \big(\|\dot{S}_0 u\|^2_{L^2} + \sum_{q\in\mathbb{N}} 2^{2qs}\|\dot{\Delta}_q u\|^2_{L^2}\big)^{1/2}$$

These are a particular case of the general nonhomogeneous Besov spaces $B^s_{p,r}$, for $s \in \mathbb{R}, p, r \in [1,\infty]^2$ consisting of all tempered distributions u such that:

$$\|u\|_{B^s_{p,r}} \overset{def}{=} \begin{cases} \|(\|\dot{S}_0 u\|^r_{L^p} + \sum_{q\in\mathbb{N}} 2^{rqs}\|\dot{\Delta}_q u\|^r_{L^p})^{\frac{1}{r}} & \text{if } r<\infty\\ \max(\|\dot{S}_0 u\|_{L^p}, \sup_{q\in\mathbb{N}} 2^{qs}\|\dot{\Delta}_q u\|_{L^p}) & \text{if } r=\infty \end{cases}$$

which reduces to the nonhomogeneous Sobolev space H^s for $p = r = 2$.

Similarly we also have the norm of the *homogenous* Sobolev spaces $\dot{H}^s$:

$$\|u\|_{\dot{H}^s} = \big(\sum_{q\in\mathbb{Z}} 2^{2qs}\|\dot{\Delta}_q u\|^2_{L^2}\big)^{1/2}$$

and the homogenous Besov spaces $\dot{B}^s_{p,r}$ for $s \in \mathbb{R}, p, r \in [1,\infty]^2$ consisting of all the homogeneous tempered distributions u such that:

$$\|u\|_{\dot{B}^s_{p,r}} \overset{def}{=} \begin{cases} \|(\sum_{q\in\mathbb{Z}} 2^{rqs}\|\dot{\Delta}_q u\|^r_{L^p})^{\frac{1}{r}} & \text{if } r<\infty\\ \sup_{q\in\mathbb{Z}} 2^{qs}\|\dot{\Delta}_q u\|_{L^p} & \text{if } r=\infty \end{cases}$$

which reduces to the homogeneous Sobolev space $\dot{H}^s$ for $p = r = 2$.

Let us note that the homogeneous Besov spaces have somewhat better product rules, and this specificity encoded in Theorem A.1 will be very useful in our subsequent estimates.

Furthermore we will need the following characterisation of the homogeneous norms, in terms of operators $\dot{S}_q u$:

Lemma 4.1. **[Prop. 2.33] , [2]** *Let $s < 0$ and $p, r \in [1,\infty]^2$. A tempered distribution u belongs to $\dot{B}^s_{p,r}$ if and only if:*

$$(2^{qs}\|\dot{S}_q u\|_{L^p})_{q\in\mathbb{Z}} \in l^r$$

and for some constant C depending only on the dimension d we have:

$$C^{-|s|+1}\|u\|_{\dot{B}^s_{p,r}} \leq \|(2^{qs}\|\dot{S}_q u\|_{L^p})_q\|_{l^r} \leq C(1+\frac{1}{|s|})\|u\|_{\dot{B}^s_{p,r}}$$

We will use the following well-known estimates:

Lemma 4.2. *([12], [13])*

(i) *(Bernstein inequalities)*

$$2^{-q}\|\nabla\dot{S}_q u\|_{L^p_x} \leq C\|u\|_{L^p_x}, \forall 1 \leq p \leq \infty$$

$$c\|\dot{\Delta}_q u\|_{L^p_x} \leq 2^{-q}\|\dot{\Delta}_q\nabla u\|_{L^p_x} \leq C\|\dot{\Delta}_q u\|_{L^p_x}, \forall 1 \leq p \leq \infty$$

(ii) *(Bernstein inequalities)*

$$\|\dot{\Delta}_q u\|_{L^b_x} \leq 2^{d(\frac{1}{a}-\frac{1}{b})q}\|\dot{\Delta}_q u\|_{L^a_x},\ \textit{for } b \geq a \geq 1$$

$$\|\dot{S}_q u\|_{L^b_x} \leq 2^{d(\frac{1}{a}-\frac{1}{b})q}\|\dot{S}_q u\|_{L^a_x},\ \textit{for } b \geq a \geq 1$$

(ii) *(commutator estimate)*

$$\|[\dot{\Delta}_q, u]v\|_{L^p_x} \leq C2^{-q}\|\nabla u\|_{L^r_x}\|v\|_{L^s_x} \tag{38}$$

with $\frac{1}{p} = \frac{1}{r} + \frac{1}{s}$. The constant C depends only on the function φ used in defining $\dot{\Delta}_q$ but not on p, r, s.

Proof. For the commutator estimate we begin by writing

$$\begin{aligned}
[\dot{\Delta}_q, u]v(x) = \dot{\Delta}_q(uv)(x) - u(x)\Delta_q v(x) &= 2^{qd}\int h(2^q y)(u(x-y)-u(x))v(x-y)\mathrm{d}y \\
&= 2^{qd}\int_{\mathbb{R}^d}\int_0^1 \frac{\partial}{\partial\tau}\Big\{h(2^q y)u(x-\tau y)v(x-y)\mathrm{d}y\Big\}\mathrm{d}\tau \\
&= -2^{qd}\int_0^1\int_{\mathbb{R}^d} h(2^q y)y\cdot\nabla u(x-\tau y)v(x-y)\mathrm{d}y\mathrm{d}\tau \\
&= -2^{-q}\int_0^1\int_{\mathbb{R}^d} \tilde{h}_{2^q}(y)\cdot\nabla u(x-\tau y)v(x-y)\mathrm{d}y\mathrm{d}\tau,
\end{aligned}$$

where $\tilde{h}(y) := yh(y) \in \mathcal{S}(R^d)^d$ and $\tilde{h}_\lambda(y) := \lambda^d\tilde{h}(\lambda y)$. Using the Cauchy-Schwartz inequality and a change of variables, we get

$$\begin{aligned}
|[\dot{\Delta}_q, u]v(x)| &\leq 2^{-q}\int_0^1 \left(\int_{\mathbb{R}^d} |\tilde{h}_{2^q}(y)||\nabla u(x-\tau y)|^{\frac{r}{p}}\mathrm{d}y\right)^{\frac{p}{r}}\mathrm{d}\tau\left(\int_{\mathbb{R}^d}|\tilde{h}_{2^q}(y)||v(x-y)|^{\frac{s}{p}}\mathrm{d}y\right)^{\frac{p}{s}} \\
&= 2^{-q}\int_0^1 \left(\int_{\mathbb{R}^d} \frac{|\tilde{h}_{2^q\tau^{-1}}(y)|}{\tau^d}|\nabla u(x-y)|^{\frac{r}{p}}\mathrm{d}y\right)^{\frac{p}{r}}\mathrm{d}\tau\left(\int_{\mathbb{R}^d}|\tilde{h}_{2^q}(y)||v(x-y)|^{\frac{s}{p}}\mathrm{d}y\right)^{\frac{p}{s}} \\
&= 2^{-q}\int_0^1 \Big(\frac{|\tilde{h}_{2^q\tau^{-1}}|}{\tau^d} * |\nabla u|^{\frac{r}{p}}(x)\Big)^{\frac{p}{r}}\mathrm{d}\tau\,\Big(|\tilde{h}_{2^q}| * |v|^{\frac{s}{p}}(x)\Big)^{\frac{p}{s}}
\end{aligned}$$

Taking the L^p norm in the x variable, using the Cauchy-Schwartz inequality in the x variable and convolution estimates we obtain

$$\begin{aligned}
\|[\dot{\Delta}_q, u]v\|_{L^p_x} &\leq 2^{-q}\int_0^1 \Big\|\Big(\frac{|\tilde{h}_{2^q\tau^{-1}}|}{\tau^d} * |\nabla u|^{\frac{r}{p}}(x)\Big)^{\frac{p}{r}}\Big\|_{L^r_x}\mathrm{d}\tau\,\Big\|\Big(|\tilde{h}_{2^q}| * |v|^{\frac{s}{p}}(x)\Big)^{\frac{p}{s}}\Big\|_{L^s_x} \\
&\leq 2^{-q}\Big(\int_0^1 \|\frac{|\tilde{h}_{2^q\tau^{-1}}|}{\tau^d} * |\nabla u|^{\frac{r}{p}}\|^{\frac{p}{r}}_{L^p_x}\mathrm{d}\tau\Big)\||\tilde{h}_{2^q}| * |v|^{\frac{s}{p}}\|^{\frac{p}{s}}_{L^p_x} \\
&\leq 2^{-q}\int_0^1 \frac{\|\tilde{h}_{2^q\tau^{-1}}\|^{\frac{p}{r}}_{L^1_x}}{\tau^d}\mathrm{d}\tau\|\nabla u\|_{L^r_x}\|\tilde{h}_{2^q}\|^{\frac{p}{s}}_{L^1_x}\|v\|_{L^s_x} \\
&\leq 2^{-q}\|\tilde{h}_{2^{-q}}\|^{\frac{p}{r}}_{L^1}\|\tilde{h}_{2^{-q}}\|^{\frac{p}{s}}_{L^1}\|\nabla u\|_{L^r_x}\|v\|_{L^s_x}.
\end{aligned}$$

Now, since

$$\|\tilde{h}_{2^{-q}}\|_{L^1_x} = \int_{\mathbb{R}^d} 2^{-qd}|\tilde{h}(2^{-q}x)|\mathrm{d}x = \int_{\mathbb{R}^d}|\tilde{h}(y)|\mathrm{d}y = \|\tilde{h}\|_{L^1_x},$$

we finally obtain

$$\|[\dot{\Delta}_q, u]v\|_{L^p_x} \leq 2^{-q}\|\tilde{h}\|^{\frac{p}{r}}_{L^1_x}\|\tilde{h}\|^{\frac{p}{s}}_{L^1}\|\nabla u\|_{L^r_x}\|v\|_{L^s_x} = \|\tilde{h}\|_{L^1_x}2^{-q}\|\nabla u\|_{L^r_x}\|v\|_{L^s_x}$$

so the constant in the inequality is $C = \|\tilde{h}\|_{L^1}$ and it does not depend on p, r, s. $\square$

We will also make use of a Bernstein-type inequality evolving the operator $\dot{S}_q$.

Lemma 4.3. *there exist two positive constants $\tilde{c}$ and $\tilde{C}$ such that*

$$\tilde{c}\|(\dot{S}_q - \dot{S}_{q'})u\|_{L^p_x} \leq 2^{-q}\|(\dot{S}_q - \dot{S}_{q'})\nabla u\|_{L^p_x} \leq \tilde{C}\|(\dot{S}_q - \dot{S}_{q'})u\|_{L^p_x}, \forall 1 \leq p \leq \infty,$$

for any integers q and q' with $|q - q'| \leq 5$.

Proof. First, we consider new localizer functions as follows:

$$\tilde{\varphi}_q(\xi) := \frac{1}{10}\sum_{|q-j|\leq 10}\varphi_j(\xi) \quad \text{and} \quad \tilde{\chi}(\xi) := \begin{cases} \sum_{q\leq -1}\tilde{\varphi}_q(\xi) & \text{if} \quad \xi \neq 0, \\ 1 & \text{otherwise,} \end{cases}$$

so that (35) and (36) are satisfied with $\tilde{\varphi}$ and $\tilde{\chi}$ instead of φ and χ. Then defining the new homogeneous dyadic block $\dot{\tilde{\Delta}}_q$ in the same line of $\dot{\Delta}_q$, we have

$$\dot{\tilde{\Delta}}_q(\dot{S}_q - \dot{S}_{q'})u = \frac{1}{10}\sum_{|q-j|\leq 10}\dot{\Delta}_j(\dot{S}_q - \dot{S}_{q'})(u) = \frac{1}{10}(\dot{S}_q - \dot{S}_{q'})(u).$$

Then the inequality turns out from **(i)** of Lemma 4.2, making use of $\dot{\tilde{\Delta}}_q$ instead of $\dot{\Delta}_q$. $\square$

4.2. The proof of the uniqueness. In this section we provide the proof of the uniqueness result for the weak solutions of system (8). The main idea is to evaluate the difference between two weak solutions in a functional space which is less regular than L^2_x such as $\dot{H}^{-\frac{1}{2}}$. Such strategy is not new in literature, for instance we recall [15] and [29]. We now provide the uniqueness part of the proof of Theorem 1.1.

Proof. Let us consider two weak solutions $(u_1,\, Q_1)$ and $(u_2,\, Q_2)$ of system (8). We denote $\delta u := u_1 - u_2$ and $\delta Q := Q_1 - Q_2$ while $\delta S(Q,\, \nabla u)$ stands for $S(Q_1,\, \nabla u_1) - S(Q_2,\, \nabla u_2)$. Similarly, we define $\delta H(Q)$, $\delta F(Q)$, $\delta\tau$ and $\delta\sigma$. Thus $(\delta u,\, \delta Q)$ is a weak solution of

$$\begin{cases} \partial_t \delta Q - L\Delta\delta Q = \delta S(Q,\nabla u) + \Gamma \delta H(Q) - \delta u\cdot\nabla Q_1 - u_2\cdot\nabla\delta Q & \mathbb{R}_+\times\mathbb{R}^2,\\ \partial_t\delta u - \Delta\delta u + \nabla\delta\Pi = L\,\mathrm{div}\{\delta\tau+\delta\sigma\} - \delta u\cdot\nabla u_1 - u_2\cdot\nabla\delta u & \mathbb{R}_+\times\mathbb{R}^2,\\ \mathrm{div}\,\delta u = 0 & \mathbb{R}_+\times\mathbb{R}^2,\\ (\delta u,\, \delta Q)_{t=0} = (0,\, 0) & \mathbb{R}^2. \end{cases} \tag{39}$$

First, let us explicitly state $\delta S(Q,\, \nabla u)$, $\delta F(Q)$, $\delta\tau$ and $\delta\sigma$ in terms of δQ and δu, namely:

$$\begin{aligned} \delta S(Q,\nabla u) = &+(\xi\delta D + \delta\Omega)\delta Q + (\xi\delta D+\delta\Omega)(\,Q_2 + \frac{\mathrm{Id}}{2}\,) + (\xi D_2 + \Omega_2)\delta Q + \delta Q(\xi\delta D - \delta\Omega) +\\ &+(\,Q_2+\frac{\mathrm{Id}}{2}\,)(\xi\delta D - \delta\Omega) + \delta Q(\xi D_2 - \Omega_2) - 2\xi\delta Q\,\mathrm{tr}(\delta Q\nabla\delta u) - 2\xi\delta Q\,\mathrm{tr}(\delta Q\nabla u_2) +\\ &- 2\xi\delta Q\,\mathrm{tr}(Q_2\nabla\delta u) - 2\xi(\,Q_2+\frac{\mathrm{Id}}{2}\,)\mathrm{tr}(\delta Q\nabla\delta u) - 2\xi\delta Q\,\mathrm{tr}(Q_2\nabla u_2) +\\ &- 2\xi(\,Q_2+\frac{\mathrm{Id}}{2}\,)\mathrm{tr}(\delta Q\nabla u_2) - 2\xi(\,Q_2+\frac{\mathrm{Id}}{2}\,)\mathrm{tr}(Q_2\nabla\delta u), \end{aligned}$$

$$\delta F(Q) = -a\,\delta Q + b\,(\,Q_1\delta Q + \delta Q Q_2\,) - b\,\mathrm{tr}\{\delta Q Q_1 + Q_2\delta Q\}\frac{\mathrm{Id}}{2} + -c\left[\delta Q\mathrm{tr}\{Q_1^2\} + Q_2\mathrm{tr}\{\delta Q Q_1 + Q_2\delta Q\}\right]$$

$$\delta H(Q) = \delta F(Q) + L\Delta\delta Q.$$

$$\begin{aligned} \delta\tau = &-\xi\delta Q\,F(Q_1) - \xi(\,Q_2+\frac{\mathrm{Id}}{2}\,)\delta F(Q) - L\xi\delta Q\,\Delta\delta Q - L\xi\delta Q\,\Delta Q_2 - L\xi(\,Q_2+\frac{\mathrm{Id}}{2}\,)\Delta\delta Q +\\ &- \xi F(Q_1)\delta Q - \xi\delta F(Q)(\,Q_2+\frac{\mathrm{Id}}{2}\,) - L\xi\Delta\delta Q\delta Q - L\xi\Delta Q_2\delta Q - L\xi\Delta\delta Q(\,Q_2+\frac{\mathrm{Id}}{2}\,) +\\ &+ 2\xi\delta Q\mathrm{tr}\{Q_1F(Q_1)\} + 2\xi Q_2\mathrm{tr}\{\delta Q F(Q_1)\} +\\ &+ 2\xi Q_2\mathrm{tr}\{Q_2\delta F(Q)\} + 2L\xi\delta Q\mathrm{tr}\{\delta Q\Delta\delta Q\} + 2L\xi\delta Q\mathrm{tr}\{\delta Q\Delta Q_2\} + 2L\xi\delta Q\mathrm{tr}\{Q_2\Delta\delta Q\} +\\ &+ 2L\xi(\,Q_2+\frac{\mathrm{Id}}{2}\,)\mathrm{tr}\{\delta Q\Delta\delta Q\} + 2L\xi\delta Q\mathrm{tr}\{Q_2\Delta Q_2\} + 2L\xi(\,Q_2+\frac{\mathrm{Id}}{2}\,)\mathrm{tr}\{\delta Q\Delta Q_2\} +\\ &+ 2L\xi(\,Q_2+\frac{\mathrm{Id}}{2}\,)\mathrm{tr}\{Q_2\Delta\delta Q\} - L\nabla\delta Q\odot\nabla Q_1 - L\nabla Q_2\odot\nabla\delta Q - L\frac{\mathrm{Id}}{2}\mathrm{tr}\{\delta Q Q_1\} - L\frac{\mathrm{Id}}{2}\mathrm{tr}\{Q_2\delta Q\} \end{aligned}$$

$$\begin{aligned} \delta\sigma = \delta Q F(Q_1) + Q_2\delta F(Q) - F(Q_1)\delta Q - \delta F(Q)Q_2 + L\delta Q\Delta\delta Q + LQ_2\Delta\delta Q + L\delta Q\Delta Q_2&\\ - L\Delta\delta Q\delta Q - L\Delta Q_2\delta Q - L\Delta\delta Q Q_2& \end{aligned}$$

Taking the inner product in $\dot{H}^{-1/2}$ of the first equation with $-L\lambda\Delta\delta Q$ and adding to it the scalar product in $\dot{H}^{-1/2}$ of the second one by $\frac{1}{\lambda}\delta u$ we get:

$$\begin{aligned} &\frac{\mathrm{d}}{\mathrm{d}t}\Big[\frac{1}{2\lambda}\|\delta u\|^2_{\dot{H}^{-\frac{1}{2}}} + L\|\nabla\delta Q\|^2_{\dot{H}^{-\frac{1}{2}}}\Big] + \frac{\nu}{\lambda}\|\nabla\delta u\|^2_{\dot{H}^{-\frac{1}{2}}} + \Gamma L^2\|\Delta\delta Q\|^2_{\dot{H}^{-\frac{1}{2}}} =\\ &- L\langle(\xi\delta D+\delta\Omega)\delta Q,\Delta\delta Q\rangle \underbrace{-L\xi\langle\delta D\,Q_2,\Delta\delta Q\rangle}_{\mathcal{A}_1}\underbrace{-L\langle\delta\Omega\,Q_2,\Delta\delta Q\rangle}_{\mathcal{B}_1}\underbrace{-L\xi\langle\frac{\delta D}{2},\Delta\delta Q\rangle}_{\mathcal{C}_1}\underbrace{-L\langle\frac{\delta\Omega}{2},\Delta\delta Q\rangle}_{\mathcal{D}_1}\\ &- L\langle(\xi D_2+\Omega_2)\delta Q,\Delta\delta Q\rangle - L\langle\delta Q(\xi\delta D-\delta\Omega),\Delta\delta Q\rangle\underbrace{-L\xi\langle Q_2\delta D,\Delta\delta Q\rangle}_{\mathcal{A}_2}\underbrace{+L\langle Q_2\delta\Omega,\Delta\delta Q\rangle}_{\mathcal{B}_2}\\ &\underbrace{-L\xi\langle\frac{\delta D}{2},\Delta\delta Q\rangle}_{\mathcal{C}_2}\underbrace{+L\langle\frac{\delta\Omega}{2},\Delta\delta Q\rangle}_{\mathcal{D}_2} - L\langle\delta Q(\xi D_2-\Omega_2),\Delta\delta Q\rangle + 2L\xi\langle\delta Q\,\mathrm{tr}(\delta Q\nabla\delta u),\Delta\delta Q\rangle\\ &+ 2L\xi\langle\delta Q\,\mathrm{tr}(\delta Q\nabla u_2),\Delta\delta Q\rangle + 2L\xi\langle\delta Q\,\mathrm{tr}(Q_2\nabla\delta u),\Delta\delta Q\rangle + 2L\xi\langle Q_2\,\mathrm{tr}(\delta Q\nabla\delta u),\Delta\delta Q\rangle\\ &\underbrace{+2L\xi\langle\frac{\mathrm{Id}}{2}\mathrm{tr}(\delta Q\nabla\delta u),\Delta\delta Q\rangle}_{=0} + 2L\xi\langle\delta Q\,\mathrm{tr}(Q_2\nabla u_2),\Delta\delta Q\rangle + 2L\xi\langle Q_2\,\mathrm{tr}(\delta Q\nabla u_2),\Delta\delta Q\rangle \end{aligned}$$

$$
\begin{aligned}
&\underbrace{+2L\xi\langle \frac{\mathrm{Id}}{2}\mathrm{tr}(\delta Q\nabla u_2),\Delta\delta Q\rangle}_{=0}\underbrace{+2L\xi\langle Q_2\,\mathrm{tr}(Q_2\nabla\delta u),\Delta\delta Q\rangle}_{\mathcal{E}_1}\underbrace{+2L\xi\langle\frac{\mathrm{Id}}{2}\mathrm{tr}(Q_2\nabla\delta u),\Delta\delta Q\rangle}_{=0}\\
&+La\Gamma\langle\delta Q,\Delta\delta Q\rangle-Lb\Gamma\langle Q_1\delta Q+\delta QQ_2,\Delta\delta Q\rangle\underbrace{+Lb\Gamma\langle\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\}\frac{\mathrm{Id}}{2},\Delta\delta Q\rangle}_{=0}\\
&+Lc\Gamma\langle\delta Q\mathrm{tr}\{Q_1^2\},\Delta\delta Q\rangle+Lc\Gamma\langle Q_2\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\},\Delta\delta Q\rangle+L\langle\delta u\cdot\nabla Q_1,\Delta\delta Q\rangle\\
&+L\langle u_2\cdot\nabla\delta Q,\Delta\delta Q\rangle-a\xi\langle\delta QQ_1,\nabla\delta u\rangle+b\xi\langle\delta QQ_1^2,\nabla\delta u\rangle-b\xi\langle\delta Q\mathrm{tr}(Q_1^2)\frac{\mathrm{Id}}{2},\nabla\delta u\rangle\\
&-c\xi\langle\delta Q\mathrm{tr}(Q_1^2)Q_1,\nabla\delta u\rangle-a\xi\langle(Q_2+\frac{\mathrm{Id}}{2})\delta Q,\nabla\delta u\rangle+b\xi\langle(Q_2+\frac{\mathrm{Id}}{2})(Q_1\delta Q+\delta QQ_2),\nabla\delta u\rangle\\
&-b\xi\langle\frac{Q_2}{2}\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\},\nabla\delta u\rangle\underbrace{-b\xi\langle\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\}\frac{\mathrm{Id}}{9},\nabla\delta u\rangle}_{=0}-c\xi\langle(Q_2+\frac{\mathrm{Id}}{2})\delta Q\mathrm{tr}\{Q_1^2\},\nabla\delta u\rangle\\
&-c\xi\langle(Q_2+\frac{\mathrm{Id}}{2})Q_2\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\},\nabla\delta u\rangle+L\xi\langle\delta Q\Delta\delta Q,\nabla\delta u\rangle+L\xi\langle\delta Q\Delta Q_2,\nabla\delta u\rangle\\
&\underbrace{+L\xi\langle Q_2\Delta\delta Q,\nabla\delta u\rangle}_{\mathcal{A}_3}\underbrace{+L\xi\langle\frac{\Delta\delta Q}{2},\nabla\delta u\rangle}_{\mathcal{C}_3}-a\xi\langle Q_1\delta Q,\nabla\delta u\rangle+b\xi\langle(Q_1^2-\mathrm{tr}\{Q_1^2\}\frac{\mathrm{Id}}{2})\delta Q,\nabla\delta u\rangle\\
&-c\xi\langle Q_1^2\mathrm{tr}\{Q_1^2\}\delta Q,\nabla\delta u\rangle-a\xi\langle\delta Q(Q_2+\frac{\mathrm{Id}}{2}),\nabla\delta u\rangle+b\xi\langle(Q_1\delta Q+\delta QQ_2)(Q_2+\frac{\mathrm{Id}}{2}),\nabla\delta u\rangle\\
&-b\xi\langle\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\}\frac{Q_2}{2},\nabla\delta u\rangle\underbrace{-b\xi\langle\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\}\frac{\mathrm{Id}}{9},\nabla\delta u\rangle}_{=0}-c\xi\langle\delta Q\mathrm{tr}\{Q_1^2\}(Q_2+\frac{\mathrm{Id}}{2}),\nabla\delta u\rangle\\
&-c\xi\langle Q_2\mathrm{tr}\{\delta Q\delta Q_1+Q_2\delta Q\}(Q_2+\frac{\mathrm{Id}}{2}),\nabla\delta u\rangle+L\xi\langle\Delta\delta Q\,\delta Q,\nabla\delta u\rangle+L\xi\langle\Delta Q_2\delta Q,\nabla\delta u\rangle\\
&\underbrace{+L\xi\langle\Delta\delta QQ_2,\nabla\delta u\rangle}_{\mathcal{A}_4}\underbrace{+L\xi\langle\frac{\Delta\delta Q}{2},\nabla\delta u\rangle}_{\mathcal{C}_4}+2a\xi\langle\delta Q\mathrm{tr}\{Q_1^2\},\nabla\delta u\rangle-2b\xi\langle\delta Q\mathrm{tr}\{Q_1^3\},\nabla\delta u\rangle\\
&\underbrace{+2b\xi\langle\frac{\delta Q}{2}\mathrm{tr}\{Q_1\}\mathrm{tr}\{Q_1^2\},\nabla\delta u\rangle}_{=0}+2c\xi\langle\delta Q\mathrm{tr}\{Q_1^2\}^2,\nabla\delta u\rangle+2a\xi\langle Q_2\mathrm{tr}\{\delta QQ_1\},\nabla\delta u\rangle\\
&-2b\xi\langle Q_2\mathrm{tr}\{\delta QQ_1^2\},\nabla\delta u\rangle\underbrace{+2b\xi\langle\frac{Q_2}{2}\mathrm{tr}\{\delta Q\}\mathrm{tr}\{Q_1^2\},\nabla\delta u\rangle}_{=0}+2c\xi\langle Q_2\mathrm{tr}\{\delta QQ_1\}\mathrm{tr}\{Q_1^2\},\nabla\delta u\rangle\\
&+2a\xi\langle Q_2\mathrm{tr}\{Q_2\delta Q\},\nabla\delta u\rangle-2b\xi\langle Q_2\mathrm{tr}\{Q_2(Q_1\delta Q+\delta QQ_2)\},\nabla\delta u\rangle\\
&\underbrace{+2b\xi\langle Q_2\mathrm{tr}\{\frac{Q_2}{2}\}\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\},\nabla\delta u\rangle}_{=0}+2c\xi\langle Q_2\mathrm{tr}\{Q_2\delta Q\}\mathrm{tr}\{Q_1^2\},\nabla\delta u\rangle\\
&+2c\xi\langle Q_2\mathrm{tr}\{Q_2^2\}\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\},\nabla\delta u\rangle-2L\xi\langle\delta Q\mathrm{tr}\{\delta Q\Delta\delta Q\},\nabla\delta u\rangle\\
&-2L\xi\langle\delta Q\mathrm{tr}\{\delta Q\Delta Q_2\},\nabla\delta u\rangle-2L\xi\langle\delta Q\mathrm{tr}\{Q_2\Delta\delta Q\},\nabla\delta u\rangle\\
&-2L\xi\langle Q_2\mathrm{tr}\{\delta Q\Delta\delta Q\},\nabla\delta u\rangle\underbrace{-2L\xi\langle\frac{\mathrm{Id}}{2}\mathrm{tr}\{\delta Q\Delta\delta Q\},\nabla\delta u\rangle}_{=0}-2L\xi\langle\delta Q\mathrm{tr}\{Q_2\Delta Q_2\},\nabla\delta u\rangle\\
&-2L\xi\langle Q_2\mathrm{tr}\{\delta Q\Delta Q_2\},\nabla\delta u\rangle\underbrace{-2L\xi\langle\frac{\mathrm{Id}}{2}\mathrm{tr}\{\delta Q\Delta Q_2\},\nabla\delta u\rangle}_{=0}\underbrace{-2L\xi\langle Q_2\mathrm{tr}\{Q_2\Delta\delta Q\},\nabla\delta u\rangle}_{\mathcal{E}_2}\\
&\underbrace{-2L\xi\langle\frac{\mathrm{Id}}{2}\mathrm{tr}\{Q_2\Delta\delta Q\},\nabla\delta u\rangle}_{=0}+L\langle\nabla\delta Q\odot\nabla Q_1,\nabla\delta u\rangle+L\langle\nabla Q_2\odot\nabla\delta Q_1,\nabla\delta u\rangle\underbrace{+L\langle\frac{\mathrm{Id}}{2}\mathrm{tr}\{\delta QQ_1\},\nabla\delta u\rangle}_{=0}\\
&\underbrace{+L\langle\frac{\mathrm{Id}}{2}\mathrm{tr}\{Q_2\delta Q\},\nabla\delta u\rangle}_{=0}+La\langle\delta QQ_1,\nabla\delta u\rangle-Lb\langle\delta Q(Q_1^2-\mathrm{tr}\{Q_1^2\}\frac{\mathrm{Id}}{2}),\nabla\delta u\rangle+Lc\langle\delta QQ_1\mathrm{tr}\{Q_1^2\},\nabla\delta u\rangle\\
&+a\langle Q_2\delta Q,\nabla\delta u\rangle-b\langle Q_2(Q_1\delta Q+\delta QQ_2),\nabla\delta u\rangle+b\langle Q_2\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\}\frac{\mathrm{Id}}{2},\nabla\delta u\rangle\\
&+c\langle Q_2\delta Q\mathrm{tr}\{Q_1^2\},\nabla\delta u\rangle\underbrace{+c\langle Q_2^2\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\},\nabla\delta u\rangle}_{\mathcal{F}_1}\\
&-a\langle Q_1\delta Q,\nabla\delta u\rangle+b\langle(Q_1^2-\mathrm{tr}\{Q_1^2\}\frac{\mathrm{Id}}{2})\delta Q,\nabla\delta u\rangle-c\langle Q_1\mathrm{tr}\{Q_1^2\}\delta Q,\nabla\delta u\rangle-a\langle\delta QQ_2,\nabla\delta u\rangle
\end{aligned}
\tag{40}
$$

$$
\begin{aligned}
&+ b\langle (Q_1\delta Q + \delta Q Q_2)Q_2, \nabla\delta u\rangle - b\langle \mathrm{tr}\{\delta Q Q_1 + Q_2\delta Q\}\frac{\mathrm{Id}}{2}Q_2, \nabla\delta u\rangle - c\langle \delta Q\,\mathrm{tr}\{Q_1^2\}Q_2, \nabla\delta u\rangle\\
&\underbrace{-c\langle Q_2\mathrm{tr}\{\delta Q Q_1 + Q_2\delta Q\}Q_2, \nabla\delta u\rangle}_{\mathcal{F}_2} - L\langle \delta Q\Delta\delta Q, \nabla\delta u\rangle \underbrace{-L\langle Q_2\Delta\delta Q,\nabla\delta u\rangle}_{\mathcal{B}_3} - L\langle\delta Q\Delta Q_2,\nabla\delta u\rangle\\
&+ L\langle\Delta\delta Q\delta Q,\nabla\delta u\rangle \underbrace{+L\langle\Delta\delta Q Q_2,\nabla\delta u\rangle}_{\mathcal{B}_4} + L\langle\Delta Q_2\delta Q,\nabla\delta u\rangle - \langle u_2\cdot\nabla\delta u,\delta u\rangle - \langle\delta u\cdot\nabla u_1,\delta u\rangle.
\end{aligned}
$$

Denoting

$$\Phi(t) := 1/(2\lambda)\|\delta u(t)\|_{\dot H^{-1/2}}^2 + L\|\nabla\delta Q(t)\|_{\dot H^{-1/2}}^2$$

we will aim to show that Φ satisfies the inequality

$$\Phi'(t) \leq \chi(t)\mu(\Phi(t)), \tag{41}$$

where μ is an Osgood modulus of continuity (see [2], Definition 3.1), given by

$$\mu(r) := r + r\ln\Big(1+e+\frac{1}{r}\Big) + r\ln\Big(1+e+\frac{1}{r}\Big)\ln\ln\Big(1+e+\frac{1}{r}\Big). \tag{42}$$

with $\chi \in L^1_{loc}$ apriori. We are going to find a double-logarithmic estimate, hence thanks to the Osgood Lemma (see [2], Lemma 3.4) and since $\Phi(0)$ is null, we get that $\Phi\equiv 0$, which yields the uniqueness of the solution for system (8).

First, let us observe following simplifications of (40):

$$0 = \mathcal{C}_1 + \mathcal{C}_2 + \mathcal{C}_3 + \mathcal{C}_4 = \mathcal{D}_1 + \mathcal{D}_2 = \mathcal{F}_1 + \mathcal{F}_2.$$

The key method we use to obtain the desired estimates is the para-differential calculus decomposition summarized in the following:

Remark 4.4. *Let q be an integer, and A, B be $d\times d$ matrices whose components are homogeneous temperate distributions. We are going to use the following notation:*

$$
\begin{aligned}
&\mathcal{J}_q^1(A,B) := \textstyle\sum_{|q-q'|\leq 5}[\dot\Delta_q,\, \dot S_{q'-1}A]\dot\Delta_{q'}B, &&\mathcal{J}_q^3(A,B) := \dot S_{q-1}A\dot\Delta_q B,\\
&\mathcal{J}_q^2(A,B) := \textstyle\sum_{|q-q'|\leq 5}(\dot S_{q'-1}A - \dot S_{q-1}A)\dot\Delta_q\dot\Delta_{q'}B, &&\mathcal{J}_q^4(A,B) := \textstyle\sum_{q'\geq q-5}\dot\Delta_q(\dot\Delta_{q'}A\,\dot S_{q'+2}B).
\end{aligned}
$$

Than we can decompose the product AB as follows

$$\dot\Delta_q(AB) = \mathcal{J}_q^1(A,B) + \mathcal{J}_q^2(A,B) + \mathcal{J}_q^3(A,B) + \mathcal{J}_q^4(A,B) \tag{43}$$

for any integer q, thanks to (37).

Moreover from now on we will use the notation $\lesssim$ as follows: for any non-negative real numbers a and b, we denote $a \lesssim b$ if and only if there exists a positive constant C (independent of a and b) such that $a\leq Cb$.

4.2.1. *Estimate of $\mathcal{A}_1 + \mathcal{A}_2 + \mathcal{A}_3 + \mathcal{A}_4$.* Let us begin analyzing the terms $\mathcal{A}_1$, $\mathcal{A}_2$, $\mathcal{A}_3$ and $\mathcal{A}_4$ of (40). First, we observe that

$$\mathcal{A}_2 = -L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\dot\Delta_q(Q_2\delta D), \dot\Delta_q\Delta\delta Q\rangle_{L^2_x} = -L\xi\sum_{q\in\mathbb{Z}}2^{-q}\sum_{i=1}^4\langle\mathcal{J}_q^i(Q_2,\,\delta D), \dot\Delta_q\Delta\delta Q\rangle_{L^2_x}$$

Now, when $i=1$, we have

$$
\begin{aligned}
2^{-q}\langle\mathcal{J}_q^1(Q_2,\,\delta D), \dot\Delta_q\Delta\delta Q\rangle_{L^2_x} &= \sum_{|q-q'|\leq 5}2^{-q}\langle[\dot\Delta_q,\,\dot S_{q'-1}Q_2]\dot\Delta_{q'}\delta D, \dot\Delta_q\Delta\delta Q\rangle_{L^2_x}\\
&\lesssim \sum_{|q-q'|\leq 5}2^{-q}\|[\dot\Delta_q,\,\dot S_{q'-1}Q_2]\dot\Delta_{q'}\delta D\|_{L^2_x}\|\dot\Delta_q\Delta\delta Q\|_{L^2_x}\\
&\lesssim \sum_{|q-q'|\leq 5}2^{-2q}\|\dot S_{q'-1}\nabla Q_2\|_{L^4_x}\|\dot\Delta_{q'}\delta D\|_{L^4_x}\|\dot\Delta_q\Delta\delta Q\|_{L^2_x}\\
&\lesssim \sum_{|q-q'|\leq 5}2^{-q}\|\dot S_{q'-1}\nabla Q_2\|_{L^2_x}^{\frac12}\|\dot S_{q'-1}\Delta Q_2\|_{L^2_x}^{\frac12}\|\dot\Delta_{q'}\delta u\|_{L^4_x}\|\dot\Delta_q\Delta\delta Q\|_{L^2_x}\\
&\lesssim \sum_{|q-q'|\leq 5}\|\nabla Q_2\|_{L^2_x}^{\frac12}\|\Delta Q_2\|_{L^2_x}^{\frac12}\|\dot\Delta_{q'}\delta u\|_{L^2_x}2^{-\frac{q}{2}}\|\dot\Delta_q\Delta\delta Q\|_{L^2_x},
\end{aligned}
\tag{44}
$$

for every $q \in \mathbb{Z}$. Hence, we get

$$\begin{aligned}
-L\xi \sum_{q\in\mathbb{Z}} 2^{-q} \langle \mathcal{J}_q^1(Q_2,\, \delta D), \dot{\Delta}_q \Delta\delta Q\rangle_{L^2_x} &\lesssim \|\nabla Q_2\|_{L^2_x}^{\frac12} \|\Delta Q_2\|_{L^2_x}^{\frac12} \|\delta u\|_{L^2_x} \|\Delta\delta Q\|_{\dot{H}^{-\frac12}} \\
&\lesssim \|\nabla Q_2\|_{L^2_x}^{\frac12} \|\Delta Q_2\|_{L^2_x}^{\frac12} \|\delta u\|_{\dot{H}^{-\frac12}}^{\frac12} \|\nabla\delta u\|_{\dot{H}^{-\frac12}}^{\frac12} \|\Delta\delta Q\|_{\dot{H}^{-\frac12}} \\
&\lesssim \|\nabla Q_2\|_{L^2_x}^2 \|\Delta Q_2\|_{L^2_x}^2 \|\delta u\|_{\dot{H}^{-\frac12}}^2 + C_\nu \|\nabla\delta u\|_{\dot{H}^{-\frac12}}^2 + C_{\Gamma,L} \|\Delta\delta Q\|_{\dot{H}^{-\frac12}}^2,
\end{aligned} \tag{45}$$

where we have used the following interpolation inequality:

$$\|\delta u\|_{L^2_x} \le \|\delta u\|_{\dot{H}^{-\frac12}}^{\frac12} \|\delta u\|_{\dot{H}^{\frac12}}^{\frac12} = \|\delta u\|_{\dot{H}^{-\frac12}}^{\frac12} \|\nabla\delta u\|_{\dot{H}^{-\frac12}}^{\frac12}.$$

When $i = 2$, the following inequalities are fulfilled:

$$\begin{aligned}
2^{-q} \langle \mathcal{J}_q^2(Q_2,\, \delta D), \dot{\Delta}_q \Delta\delta Q\rangle_{L^2_x} &= \sum_{|q-q'|\le 5} 2^{-q} \langle (\dot{S}_{q'-1}Q_2 - \dot{S}_{q-1}Q_2) \dot{\Delta}_q \dot{\Delta}_{q'} \delta D, \dot{\Delta}_q \Delta\delta Q\rangle_{L^2_x} \\
&\lesssim \sum_{|q-q'|\le 5} 2^{-q} \|(\dot{S}_{q'-1}Q_2 - \dot{S}_{q-1}Q_2)\|_{L^\infty_x} \|\dot{\Delta}_q \dot{\Delta}_{q'} \delta D\|_{L^2_x} \|\dot{\Delta}_q \Delta\delta Q\|_{L^2_x} \\
&\lesssim \sum_{|q-q'|\le 5} 2^{-2q} \|(\dot{S}_{q'-1}\Delta Q_2 - \dot{S}_{q-1}\Delta Q_2)\|_{L^2_x} \|\dot{\Delta}_q \dot{\Delta}_{q'} \delta D\|_{L^2_x} \|\dot{\Delta}_q \Delta\delta Q\|_{L^2_x} \\
&\lesssim \sum_{|q-q'|\le 5} 2^{-2q} \|\Delta Q_2\|_{L^2_x} \|\dot{\Delta}_{q'} \delta D\|_{L^2_x} \|\dot{\Delta}_q \Delta\delta Q\|_{L^2_x} \\
&\lesssim \sum_{|q-q'|\le 5} 2^{-\frac{q'}{2}} \|\dot{\Delta}_{q'} \delta u\|_{L^2_x} 2^{-\frac{q}{2}} \|\dot{\Delta}_q \Delta\delta Q\|_{L^2_x} \|\Delta Q_2\|_{L^2_x},
\end{aligned} \tag{46}$$

for any $q \in \mathbb{Z}$. Thus, it turns out that

$$-L\xi \sum_{q\in\mathbb{Z}} 2^{-q} \langle \mathcal{J}_q^2(Q_2,\, \delta D), \dot{\Delta}_q \Delta\delta Q\rangle_{L^2_x} \lesssim \|\Delta Q_2\|_{L^2_x}^2 \|\delta u\|_{\dot{H}^{-\frac12}}^2 + C_{\Gamma,L} \|\Delta\delta Q\|_{\dot{H}^{-\frac12}}^2 \tag{47}$$

The term corresponding to $i = 3$ cannot be estimated as before. We will see that this challenging term will be simplified. Finally, when $i = 4$, we have

$$\begin{aligned}
2^{-q} \langle \mathcal{J}_q^4(Q_2,\, \delta D), \dot{\Delta}_q \Delta\delta Q\rangle_{L^2_x} &= L 2^{-q} \sum_{q-q'\le 5} \langle \dot{\Delta}_q \big[\dot{\Delta}_{q'} Q_2 \dot{S}_{q'+2} \delta D\big],\, \dot{\Delta}_q \Delta\delta Q\rangle_{L^2_x} \\
&\lesssim 2^{-q} \sum_{q-q'\le 5} \|\dot{\Delta}_{q'} Q_2\|_{L^\infty_x} \|\dot{S}_{q'+2}\delta D\|_{L^2_x} \|\dot{\Delta}_q \Delta\delta Q\|_{L^2_x} \\
&\lesssim 2^{-q} \sum_{q-q'\le 5} 2^{-q'} \|\dot{\Delta}_{q'} \Delta Q_2\|_{L^2_x} \|\dot{S}_{q'+2}\delta D\|_{L^2_x} 2^q \|\dot{\Delta}_q \nabla\delta Q\|_{L^2_x} \\
&\lesssim \sum_{q-q'\le 5} 2^{\frac{q-q'}{2}} \|\dot{\Delta}_{q'} \Delta Q_2\|_{L^2_x} 2^{-\frac{q'+2}{2}} \|\dot{S}_{q'+2}\delta D\|_{L^2_x} 2^{-\frac{q}{2}} \|\dot{\Delta}_q \nabla\delta Q\|_{L^2_x} \\
&\lesssim \|\Delta Q_2\|_{L^2_x} 2^{-\frac{q}{2}} \|\dot{\Delta}_q \nabla\delta Q\|_{L^2_x} \sum_{q-q'\le 5} 2^{\frac{q-q'}{2}} 2^{-\frac{q'+2}{2}} \|\dot{S}_{q'+2}\delta D\|_{L^2_x},
\end{aligned} \tag{48}$$

for any $q \in \mathbb{Z}$. Hence,

$$\begin{aligned}
&-L\xi \sum_{q\in\mathbb{Z}} 2^{-q} \langle \mathcal{J}_q^4(Q_2,\, \delta D), \dot{\Delta}_q \Delta\delta Q\rangle_{L^2_x} \\
&\quad\lesssim \|\Delta Q_2\|_{L^2_x} \sum_{q\in\mathbb{Z}} 2^{-\frac{q}{2}} \|\dot{\Delta}_q \nabla\delta Q\|_{L^2_x} \sum_{q'\in\mathbb{Z}} 2^{\frac{q-q'}{2}} 1_{(-\infty,5]}(q-q') 2^{-\frac{q'+2}{2}} \|\dot{S}_{q'+2}\delta D\|_{L^2_x} \\
&\quad\lesssim \|\Delta Q_2\|_{L^2_x} \|\nabla\delta Q\|_{\dot{H}^{-\frac12}} \Big(\sum_{q\in\mathbb{Z}} \Big| \sum_{q'\in\mathbb{Z}} 2^{q-q'} 1_{(-\infty,5]}(q-q') 2^{-\frac{q'+2}{2}} \|\dot{S}_{q'+2}\delta D\|_{L^2_x} \Big|^2 \Big)^{\frac12},
\end{aligned}$$

and by convolution

$$\begin{aligned}
&\Big(\sum_{q\in\mathbb{Z}} \Big| \sum_{q'\in\mathbb{Z}} 2^{q-q'} 1_{(-\infty,5]}(q-q') 2^{-\frac{q'+2}{2}} \|\dot{S}_{q'+2}\delta D\|_{L^2_x} \Big|^2 \Big)^{\frac12} \\
&\qquad\lesssim \Big(\sum_{q\le 5} 2^q\Big) \Big(\Big(\sum_{q\in\mathbb{Z}} 2^{-q} \|\dot{S}_q \delta D\|_{L^2_x}^2 \Big)^{\frac12} \Big) \lesssim \|\nabla\delta u\|_{\dot{H}^{-\frac12}},
\end{aligned}$$

so that

$$\begin{aligned}-L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\mathcal{J}_q^4(Q_2,\,\delta D),\dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x}&\lesssim\|\Delta Q_2\|_{L^2_x}\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\&\lesssim\|\Delta Q_2\|_{L^2_x}^2\|\nabla\delta Q\|_{\dot H^{-\frac12}}^2+C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2.\end{aligned}\tag{49}$$

Summarizing, it remains to control

$$\mathcal{A}_1+\mathcal{A}_3+\mathcal{A}_4-L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\mathcal{J}_q^3(Q_2,\,\delta D),\dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x}.$$

Now, observing that

$$\begin{aligned}\mathcal{A}_1=-L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\dot{\Delta}_q(\delta DQ_2),\dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x}&=-L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\dot{\Delta}_q{}^{\rm t}(\delta DQ_2),\dot{\Delta}_q{}^{\rm t}\Delta\delta Q\rangle_{L^2_x}\\&=-L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\dot{\Delta}_q(Q_2\delta D),\dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x}=\mathcal{A}_2,\end{aligned}$$

we estimate $\mathcal{A}_1$ with the previous inequalities, so that it remains to control

$$\mathcal{A}_3+\mathcal{A}_4-2L\xi\sum_{q\in\mathbb{Z}}2^{-q}\mathcal{J}_q^3(Q_2,\,\delta D)=\mathcal{A}_3+\mathcal{A}_4-2L\xi\sum_{q\in\mathbb{Z}}2^{-q}\int_{\mathbb{R}^2}{\rm tr}\big\{\dot S_{q-1}Q_2\,\dot{\Delta}_q\delta D\dot{\Delta}_q\Delta\delta Q\big\}.$$

Now, let us consider $\mathcal{A}_3=L\xi\langle Q_2\Delta\delta Q,\nabla\delta u\rangle$. We proceed along the lines used before, namely we use the decomposition given by (43):

$$\mathcal{A}_3=L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\dot{\Delta}_q(Q_2\Delta\delta Q),\dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}=L\xi\sum_{q\in\mathbb{Z}}2^{-q}\sum_{i=1}^4\langle\mathcal{J}_q^i(Q_2,\,\Delta\delta Q),\dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}.$$

When $i=1$, proceeding as for (44), we have

$$2^{-q}\langle\mathcal{J}_q^1(Q_2,\,\Delta\delta Q),\dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}\lesssim\|\nabla Q_2\|_{L^2_x}^{\frac12}\|\Delta Q_2\|_{L^2_x}^{\frac12}\sum_{|q-q'|\le5}\|\dot{\Delta}_{q'}\nabla\delta Q\|_{L^2_x}2^{-\frac q2}\|\dot{\Delta}_q\nabla\delta u\|_{L^2_x},$$

thus, considering the sum over $q\in\mathbb{Z}$ as in (45), we deduce that

$$L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\mathcal{J}_q^1(Q_2,\,\Delta\delta Q),\dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}\lesssim\|\nabla Q_2\|_{L^2_x}^2\|\Delta Q_2\|_{L^2_x}^2\|\nabla\delta Q\|_{\dot H^{-\frac12}}^2+C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2+C_{\Gamma,L}\|\Delta\delta Q\|_{\dot H^{-\frac12}}^2.\tag{50}$$

Proceeding as for proving (46), when $i=2$, we get

$$2^{-q}\langle\mathcal{J}_q^2(Q_2,\,\Delta\delta Q),\dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}\lesssim\sum_{|q-q'|\le5}2^{-\frac q2}\|\dot{\Delta}_{q'}\delta u\|_{L^2_x}2^{-\frac q2}\|\dot{\Delta}_q\Delta\delta Q\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}$$

for every $q\in\mathbb{Z}$. Thus, as in (47), it turns out that

$$L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\mathcal{J}_q^2(Q_2,\,\Delta\delta Q),\dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}\lesssim\|\Delta Q_2\|_{L^2_x}^2\|\delta u\|_{\dot H^{-\frac12}}^2+C_{\Gamma,L}\|\Delta\delta Q\|_{\dot H^{-\frac12}}^2\tag{51}$$

Finally, with the same strategy as for (48), we observe that

$$2^{-q}\langle\mathcal{J}_q^4(Q_2,\,\Delta\delta Q),\dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}\lesssim\|\Delta Q_2\|_{L^2_x}2^{-q}\|\dot{\Delta}_q\delta u\|_{L^2_x}^2\sum_{q-q'\le5}2^{\frac{q-q'}{2}}2^{-\frac{q'+2}{2}}\|\dot S_{q'+2}\Delta\delta Q\|_{L^2_x},$$

hence, as for (49), we obtain

$$\begin{aligned}L\xi\sum_{q\in\mathbb{Z}}2^{-q}\langle\mathcal{J}_q^4(Q_2,\,\Delta\delta Q),\dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}&\lesssim\|\Delta Q_2\|_{L^2_x}\|\delta u\|_{\dot H^{-\frac12}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\&\lesssim\|\Delta Q_2\|_{L^2_x}^2\|\delta u\|_{\dot H^{-\frac12}}^2+C_{\Gamma,L}\|\Delta\delta Q\|_{\dot H^{-\frac12}}^2.\end{aligned}\tag{52}$$

Summarizing all the previous considerations, we note that it remains to control

$$\begin{aligned}&\mathcal{A}_4+L\xi\sum_{q\in\mathbb{Z}}2^{-q}\Big[\langle\mathcal{J}_q^3(Q_2,\,\Delta\delta Q),\dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}-2\int_{\mathbb{R}^2}{\rm tr}\big\{\dot S_{q-1}Q_2\,\dot{\Delta}_q\delta D\dot{\Delta}_q\Delta\delta Q\big\}\Big]=\\&=\mathcal{A}_4+L\xi\sum_{q\in\mathbb{Z}}2^{-q}\int_{\mathbb{R}^2}\Big[{\rm tr}\{\dot S_{q-1}Q_2\,\dot{\Delta}_q\Delta\delta Q\dot{\Delta}_q\nabla\delta u\}-2{\rm tr}\big\{\dot S_{q-1}Q_2\,\dot{\Delta}_q\delta D\dot{\Delta}_q\Delta\delta Q\big\}\Big].\end{aligned}$$

We handle the last term $\mathcal{A}_4$ arguing as for $\mathcal{A}_3$, since $\mathcal{A}_4$ is given by

$$L\xi\langle\dot{\Delta}_q(\Delta\delta Q Q_2), \dot{\Delta}_q\nabla\delta u\rangle_{\dot{H}^{-\frac{1}{2}}} = L\xi\langle\dot{\Delta}_q(Q_2\Delta\delta Q), {}^{\mathrm{t}}\dot{\Delta}_q\nabla\delta u\rangle_{\dot{H}^{-\frac{1}{2}}} = L\xi\sum_{q\in\mathbb{Z}}2^{-q}\sum_{i=1}^{4}\langle\mathcal{J}_q^i(Q_2,\, \Delta\delta Q),\, \dot{\Delta}_q{}^{\mathrm{t}}\nabla\delta u\rangle_{L^2_x}.$$

The terms related to $i = 1, 2, 4$ are estimated similarily as $\mathcal{A}_3$. Hence it remains to evaluate

$$L\xi\sum_{q\in\mathbb{Z}}2^{-q}\Big\{\langle\mathcal{J}_q^3(Q_2,\, \Delta\delta Q),\, \dot{\Delta}_q{}^{\mathrm{t}}\nabla\delta u\rangle_{L^2_x} + \int_{\mathbb{R}^2}\big[\mathrm{tr}\{\dot{S}_{q-1}Q_2\,\dot{\Delta}_q\Delta\delta Q\dot{\Delta}_q\nabla\delta u\} - 2\mathrm{tr}\big\{\dot{S}_{q-1}Q_2\,\dot{\Delta}_q\delta D\dot{\Delta}_q\Delta\delta Q\big\}\big]\Big\} =$$

$$= 2L\xi\sum_{q\in\mathbb{Z}}2^{-q}\int_{\mathbb{R}^2}\Big[\mathrm{tr}\{\dot{S}_{q-1}Q_2\,\dot{\Delta}_q\Delta\delta Q\dot{\Delta}_q\delta D\} - \mathrm{tr}\big\{\dot{S}_{q-1}Q_2\,\dot{\Delta}_q\delta D\dot{\Delta}_q\Delta\delta Q\big\}\Big] = 0,$$

which is a null series since the trace acts on symmetric matrices, so that we can permute their order.

4.2.2. *Estimate of* $\mathcal{B}_1+\mathcal{B}_2+\mathcal{B}_3+\mathcal{B}_4$. Now we want to estimate $\mathcal{B}_1+\mathcal{B}_2+\mathcal{B}_3+\mathcal{B}_4$, namely

$$-L\langle\delta\Omega Q_2 - Q_2\delta\Omega, \Delta\delta Q\rangle_{\dot{H}^{-\frac{1}{2}}} - L\langle Q_2\Delta\delta Q - \Delta\delta Q Q_2, \nabla\delta u\rangle_{\dot{H}^{-\frac{1}{2}}}.$$

First let us consider

$$\mathcal{B}_2 = L\langle Q_2\delta\Omega, \Delta\delta Q\rangle_{\dot{H}^{-\frac{1}{2}}} = L\sum_{q\in\mathbb{Z}}2^{-q}\langle\dot{\Delta}_q(\delta\Omega Q_2), \dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x} = L\sum_{q\in\mathbb{Z}}2^{-q}\sum_{i=1}^{4}\langle\mathcal{J}_q^i(Q_2,\, \delta\Omega),\, \dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x}.$$

Proceeding exactly as for proving (45), (47) and (49), with $\delta\Omega$ instead of δD, the following estimates are obtained:

$$L\sum_{q\in\mathbb{Z}}2^{-q}\langle\mathcal{J}_q^1(Q_2,\, \delta\Omega),\, \dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x} \lesssim \|\nabla Q_2\|_{L^2_x}^2\|\Delta Q_2\|_{L^2_x}^2\|\delta u\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_\nu\|\nabla\delta u\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_{\Gamma,L}\|\Delta\delta Q\|_{\dot{H}^{-\frac{1}{2}}}^2,$$

$$L\sum_{q\in\mathbb{Z}}2^{-q}\langle\mathcal{J}_q^2(Q_2,\, \delta\Omega),\, \dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x} \lesssim \|\Delta Q_2\|_{L^2_x}^2\|\delta u\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_{\Gamma,L}\|\Delta\delta Q\|_{\dot{H}^{-\frac{1}{2}}}^2,$$

and

$$L\sum_{q\in\mathbb{Z}}2^{-q}\langle\mathcal{J}_q^4(Q_2,\, \delta\Omega),\, \dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x} \lesssim \|\Delta Q_2\|_{L^2_x}^2\|\nabla\delta Q\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_\nu\|\nabla\delta u\|_{\dot{H}^{-\frac{1}{2}}}^2.$$

Now observing that

$$\mathcal{B}_1 = -L\langle\delta\Omega\, Q_2, \Delta\delta Q\rangle_{\dot{H}^{-\frac{1}{2}}} = -L\langle{}^{\mathrm{t}}(\delta\Omega\, Q_2), {}^{\mathrm{t}}\Delta\delta Q\rangle_{\dot{H}^{-\frac{1}{2}}} = L\langle Q_2\delta\Omega, \Delta\delta Q\rangle_{\dot{H}^{-\frac{1}{2}}} = \mathcal{B}_2,$$

it remains to control

$$\mathcal{B}_3 + \mathcal{B}_4 + 2L\sum_{q\in\mathbb{Z}}2^{-q}\langle\mathcal{J}_q^3(Q_2, \delta\Omega),\, \dot{\Delta}_q\Delta\delta Q\rangle_{L^2_x} = \mathcal{B}_3 + \mathcal{B}_4 + 2L\sum_{q\in\mathbb{Z}}2^{-q}\int_{\mathbb{R}^2}\mathrm{tr}\{\dot{S}_{q-1}Q_2\dot{\Delta}_q\delta\Omega\dot{\Delta}_q\Delta\delta Q\}.$$

Now, we turn to $\mathcal{B}_3$:

$$-\mathcal{B}_3 = L\langle Q_2\Delta\delta Q, \nabla\delta u\rangle_{\dot{H}^{-\frac{1}{2}}} = L\sum_{q\in\mathbb{Z}}2^{-q}\langle\dot{\Delta}_q(Q_2\Delta\delta Q), \dot{\Delta}_q\nabla\delta u\rangle_{L^2_x} = L\sum_{q\in\mathbb{Z}}2^{-q}\sum_{i=1}^{4}\langle\mathcal{J}_q^i(Q_2,\, \Delta\delta Q),\, \dot{\Delta}_q\nabla\delta u\rangle_{L^2_x}.$$

We remark that $\mathcal{B}_3 = -\mathcal{A}_3/\xi$, hence the terms related to $i = 1, 2, 4$ are estimated as in (50), (51) and (52). Thus it remains to control

$$\mathcal{B}_4 + L\sum_{q\in\mathbb{Z}}2^{-q}\big[\langle\mathcal{J}_q^3(Q_2,\, \Delta\delta Q),\, \dot{\Delta}_q\nabla\delta u\rangle_{L^2_x} + 2\int_{\mathbb{R}^2}\mathrm{tr}\{\dot{S}_{q-1}Q_2\dot{\Delta}_q\delta\Omega\dot{\Delta}_q\Delta\delta Q\}\big] =$$

$$= \mathcal{B}_4 + L\sum_{q\in\mathbb{Z}}2^{-q}\int_{\mathbb{R}^2}\big[\mathrm{tr}\{\dot{S}_{q-1}Q_2\dot{\Delta}_q\Delta\delta Q\dot{\Delta}_q\nabla\delta u\} + 2\mathrm{tr}\{\dot{S}_{q-1}Q_2\dot{\Delta}_q\delta\Omega\dot{\Delta}_q\Delta\delta Q\}\big].$$

Observing that $\mathcal{B}_4 = -\mathcal{A}_4/\xi$ we argue as for $\mathcal{B}_3$, hence it remains to evaluate

$$L\sum_{q\in\mathbb{Z}}2^{-q}\Big\{\langle\mathcal{J}_q^3(Q_2,\, \Delta\delta Q)\dot{\Delta}_q{}^{\mathrm{t}}\nabla\delta u\rangle_{L^2_x} + \int_{\mathbb{R}^2}\big[\mathrm{tr}\{\dot{S}_{q-1}Q_2\dot{\Delta}_q\Delta\delta Q\dot{\Delta}_q\nabla\delta u\} + 2\mathrm{tr}\{\dot{S}_{q-1}Q_2\dot{\Delta}_q\delta\Omega\dot{\Delta}_q\Delta\delta Q\}\big]\Big\} =$$

$$= 2L\sum_{q\in\mathbb{Z}}2^{-q}\int_{\mathbb{R}^2}\big[\mathrm{tr}\{\dot{S}_{q-1}Q_2\dot{\Delta}_q\Delta\delta Q\dot{\Delta}_q\delta\Omega\} + \mathrm{tr}\{\dot{S}_{q-1}Q_2\dot{\Delta}_q\delta\Omega\dot{\Delta}_q\Delta\delta Q\}\big] = 0,$$

where for the cancellation we used that $\dot{S}_{q-1}Q_2$ and $\dot{\Delta}_q\Delta\delta Q$ are symmetric while $\dot{\Delta}_q\delta\Omega$ is skew-symmetric.

4.2.3. *One-logarithmic Estimates.* In this subsection, we evaluate the terms of (40) which are related to the single-logarithmic term of the equality (41).

Estimate of $\langle \delta Q \mathrm{tr}\{Q_2 \nabla u_2\}, \Delta \delta Q\rangle$. Let us fix a positive real number $N > 0$ and split the considered term into two parts, the high and the low frequencies:

$$\langle \delta Q \mathrm{tr}\{Q_2 \nabla u_2\}, \Delta \delta Q\rangle_{\dot H^{-\frac12}} = \langle \delta Q \mathrm{tr}\{(\dot S_N Q_2)\nabla u_2\}, \Delta \delta Q\rangle_{\dot H^{-\frac12}} + \langle \delta Q \mathrm{tr}\{(\sum_{q\geq N}\dot\Delta_q Q_2)\nabla u_2\}, \Delta \delta Q\rangle_{\dot H^{-\frac12}}.$$

At first we deal with the low frequencies, observing that

$$\begin{aligned}\langle \delta Q \mathrm{tr}\{(\dot S_N Q_2)\nabla u_2\}, \Delta \delta Q\rangle_{\dot H^{-\frac12}} &\lesssim \|\delta Q \mathrm{tr}\{(\dot S_N Q_2)\nabla u_2\}\|_{\dot H^{-\frac12}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\ &\lesssim \|\delta Q\|_{\dot H^{\frac12}}\|(\dot S_N Q_2)\nabla u_2\|_{L^2_x}\|\Delta \delta Q\|_{\dot H^{-\frac12}} \lesssim \|\nabla\delta Q\|_{\dot H^{-\frac12}}\|\dot S_N Q_2\|_{L^\infty_x}\|\nabla u_2\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}},\end{aligned}$$

hence, by Theorem A.2, we get

$$\begin{aligned}\langle \delta Q \mathrm{tr}\{(\dot S_N Q_2)\nabla u_2\}, \Delta \delta Q\rangle_{\dot H^{-\frac12}} &\lesssim \|\nabla \delta Q\|_{\dot H^{-\frac12}}(\|Q_2\|_{L^2_x} + \sqrt{N}\|\nabla Q_2\|_{L^2_x})\|\nabla u_2\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\ &\lesssim (1+N)\|\nabla\delta Q\|^2_{\dot H^{-\frac12}}\|(Q_2,\, \nabla Q_2)\|^2_{L^2_x}\|\nabla u_2\|^2_{L^2_x} + C_\Gamma\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}.\end{aligned}$$

For the high frequencies, we proceed as follows:

$$\begin{aligned}\langle \delta Q \mathrm{tr}\{(\sum_{q\geq N}\dot\Delta_q Q_2)\nabla u_2\}, \Delta \delta Q\rangle_{\dot H^{-\frac12}} &\lesssim \|\delta Q \mathrm{tr}\{(\sum_{q\geq N}\dot\Delta_q Q_2)\nabla u_2\}\|_{\dot H^{-\frac12}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\ &\lesssim \|\delta Q\|_{\dot H^{\frac34}}\|(\sum_{q\geq N}\dot\Delta_q Q_2)\nabla u_2\|_{\dot H^{-\frac14}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\ &\lesssim \|(Q_1,\, Q_2)\|^{\frac14}_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^{\frac34}_{L^2_x}\|\sum_{q\geq N}\dot\Delta_q Q_2\|_{\dot H^{\frac34}}\|\nabla u_2\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\ &\lesssim \|(Q_1,\, Q_2)\|^{\frac14}_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^{\frac34}_{L^2_x}(\sum_{q\geq N}2^{\frac34 q}\|\dot\Delta_q Q_2\|_{L^2_x})\|\nabla u_2\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\ &\lesssim \|(Q_1,\, Q_2)\|^{\frac14}_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^{\frac34}_{L^2_x}(\sum_{q\geq N}2^{-\frac{q}{4}}\|\dot\Delta_q \nabla Q_2\|_{L^2_x})\|\nabla u_2\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\ &\lesssim \|(Q_1,\, Q_2)\|^{\frac14}_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^{\frac34}_{L^2_x}(\sum_{q\geq N}2^{-\frac{q}{4}})\|\nabla Q_2\|_{L^2_x}\|\nabla u_2\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\ &\lesssim 2^{-\frac{N}{4}}\|(Q_1,\, Q_2)\|^{\frac14}_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^{\frac34}_{L^2_x}\|\nabla Q_2\|_{L^2_x}\|\nabla u_2\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}}.\end{aligned}$$

Now, fixing $t > 0$ arbitrary, and taking $N = N(t) := \lceil \ln(1 + e + 1/\Phi(t))\rceil > 0$, where $\lceil\cdot\rceil$ is the ceiling function, we get

$$\begin{aligned}\langle \delta Q(t) \mathrm{tr}\{Q_2(t)\nabla u_2(t)\}, \Delta \delta Q(t)\rangle_{\dot H^{-\frac12}} &\lesssim \|(Q_2,\, \nabla Q_2)(t)\|^2_{L^2_x}\|\nabla u_2(t)\|^2_{L^2_x}\Phi(t)\ln\Big(1+e+\frac{1}{\Phi(t)}\Big)+\\ &+\|(Q_1,\, Q_2)(t)\|^{\frac12}_{L^2_x}\|\nabla(Q_1,\, Q_2)(t)\|^{\frac32}_{L^2_x}\|\nabla Q_2(t)\|^2_{L^2_x}\|\nabla u_2(t)\|^2_{L^2_x}\Phi(t) + C_\Gamma\|\Delta\delta Q(t)\|^2_{\dot H^{-\frac12}}.\end{aligned}$$

Thus we have obtained a one-logarithmic term of (41). Similarly, we can handle the estimate of the following elements:

$$\begin{aligned}&+2L\xi\langle \delta Q\,\mathrm{tr}(\delta Q\nabla\delta u), \Delta\delta Q\rangle_{\dot H^{-\frac12}} + 2L\xi\langle \delta Q\,\mathrm{tr}(\delta Q\nabla u_2), \Delta\delta Q\rangle_{\dot H^{-\frac12}} + 2L\xi\langle \delta Q\,\mathrm{tr}(Q_2\nabla \delta u), \Delta\delta Q\rangle_{\dot H^{-\frac12}}+\\ &+2L\xi\langle Q_2\,\mathrm{tr}(\delta Q\nabla\delta u), \Delta\delta Q\rangle_{\dot H^{-\frac12}} + 2L\xi\langle Q_2\,\mathrm{tr}(\delta Q\nabla u_2), \Delta\delta Q\rangle_{\dot H^{-\frac12}} - 2L\xi\langle \delta Q\mathrm{tr}\{\delta Q\Delta\delta Q\}, \nabla\delta u\rangle_{\dot H^{-\frac12}}-\\ &-2L\xi\langle \delta Q\mathrm{tr}\{\delta Q\Delta Q_2\}, \nabla\delta u\rangle_{\dot H^{-\frac12}} - 2L\xi\langle \delta Q\mathrm{tr}\{Q_2\Delta \delta Q\}, \nabla\delta u\rangle_{\dot H^{-\frac12}} - 2L\xi\langle Q_2\mathrm{tr}\{\delta Q\Delta\delta Q\}, \nabla\delta u\rangle_{\dot H^{-\frac12}}-\\ &\qquad\qquad -2L\xi\langle \delta Q\mathrm{tr}\{Q_2\Delta Q_2\}, \nabla\delta u\rangle_{\dot H^{-\frac12}} - 2L\xi\langle Q_2\mathrm{tr}\{\delta Q\Delta Q_2\}, \nabla\delta u\rangle_{\dot H^{-\frac12}}.\end{aligned}$$

4.2.4. *Double-Logarithmic Estimates.* In this subsection, we perform the most challenging estimate. Now, we want to control $\mathcal{E}_1 + \mathcal{E}_2$, namely

$$\begin{aligned}\mathcal{E}_1 + \mathcal{E}_2 &= 2L\xi\big(\langle Q_2\mathrm{tr}\{Q_2\nabla\delta u\},\, \Delta\delta Q\rangle_{\dot H^{-\frac12}} - \langle Q_2\mathrm{tr}\{Q_2\Delta\delta Q\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}}\big)\\ &= 2L\xi\sum_{q\in\mathbb{Z}}2^{-q}\int_{\mathbb{R}^2}\mathrm{tr}\big\{\,\dot\Delta_q(Q_2\mathrm{tr}\{Q_2\nabla\delta u\})\,\dot\Delta_q\Delta\delta Q - \dot\Delta_q(Q_2\mathrm{tr}\{Q_2\Delta\delta Q\})\,\dot\Delta_q\nabla\delta u\,\big\}\\ &= 2L\xi\sum_{i=1}^{4}\sum_{q\in\mathbb{Z}}2^{-q}\int_{\mathbb{R}^2}\mathrm{tr}\big\{\,\mathcal{J}^i_q(Q_2,\, \mathrm{tr}\{Q_2\nabla\delta u\}\,\mathrm{Id})\,\dot\Delta_q\Delta\delta Q - \mathcal{J}^i_q(Q_2,\, \mathrm{tr}\{Q_2\Delta\delta Q\}\,\mathrm{Id})\,\dot\Delta_q\nabla\delta u\,\big\}.\end{aligned} \tag{53}$$

We we will see that there are elements inside this decomposition that generate the double-logarithmic term in (41). We proceed by considering the indexes $i = 1, 2, 3, 4$, step by step.

Estimate of $\mathcal{J}_q^1$. We start with the term of (53) related to $i = 1$, passing trough the following decomposition:

$$\begin{aligned}\sum_{j=1}^{4} \sum_{|q-q'|\leq 5} \int_{\mathbb{R}_2} \mathrm{tr}\big\{ \big([\dot{\Delta}_q,\, \dot{S}_{q'-1}Q_2]\mathrm{tr}\{\mathcal{J}_{q'}^j(Q_2,\, \nabla\delta u)\}\,\mathrm{Id}\,\big)\, \dot{\Delta}_q \Delta\delta Q - \\ - \big([\dot{\Delta}_q,\, \dot{S}_{q'-1}Q_2]\mathrm{tr}\{\mathcal{J}_{q'}^j(Q_2,\, \Delta\delta Q)\}\,\mathrm{Id}\,\big)\, \dot{\Delta}_q \nabla\delta u\,\big\}.\end{aligned} \tag{54}$$

When $j = 1$, we have

$$\begin{aligned}\mathcal{I}_1^1(q,q',q'') := \int_{\mathbb{R}_2} \Big\{\Big([\dot{\Delta}_q,\, \dot{S}_{q'-1}Q_2]\mathrm{tr}\{[\dot{\Delta}_{q'}, \dot{S}_{q''-1}Q_2]\dot{\Delta}_{q''}\nabla\delta u)\}\,\mathrm{Id}\Big)\, \dot{\Delta}_q\Delta\delta Q + \\ - \Big([\dot{\Delta}_q,\, \dot{S}_{q'-1}Q_2]\mathrm{tr}\{[\dot{\Delta}_{q'}, \dot{S}_{q''-1}Q_2]\dot{\Delta}_{q''}\Delta\delta Q)\}\,\mathrm{Id}\Big)\dot{\Delta}_q\nabla\delta u\Big\} \\ \lesssim 2^{-q}\|\dot{S}_{q'-1}\nabla Q_2\|_{L^\infty_x} 2^{-q'}\|\dot{S}_{q''-1}\nabla Q_2\|_{L^\infty_x}\|\dot{\Delta}_{q''}(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}\|\dot{\Delta}_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ \lesssim 2^{-q-q'}2^{\frac{q'}{2}}\|\dot{S}_{q'-1}\nabla Q_2\|_{L^4_x}2^{\frac{q''}{2}}\|\dot{S}_{q''-1}\nabla Q_2\|_{L^4_x}2^{q''}\|\dot{\Delta}_{q''}(\delta u,\, \nabla\delta Q)\|_{L^2_x}2^q\|\dot{\Delta}_q(\delta u,\, \nabla\delta Q)\|_{L^2_x},\end{aligned}$$

which yields

$$\mathcal{I}_1^1(q,q',q'') \lesssim 2^{\frac{3q''}{2}-\frac{q'}{2}}\|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|\dot{\Delta}_{q''}(\delta u,\, \nabla\delta Q)\|_{L^2_x}\|\dot{\Delta}_q(\delta u,\, \nabla\delta Q)\|_{L^2_x}. \tag{55}$$

Hence, taking the sum, we deduce that

$$\begin{aligned}2L\xi\sum_{q\in\mathbb{Z}}\sum_{|q-q'|\leq 5}\sum_{|q'-q''|\leq 5} 2^{-q}\mathcal{I}_1^1(q,q',q'') \lesssim \\ \lesssim \sum_{q\in\mathbb{Z}}\sum_{|q-q'|\leq 5}\sum_{|q'-q''|\leq 5} 2^{-q}2^{\frac{3q''}{2}-\frac{q'}{2}}\|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|\dot{\Delta}_{q''}(\delta u,\, \nabla\delta Q)\|_{L^2_x}\|\dot{\Delta}_q(\delta u,\, \nabla\delta Q)\|_{L^2_x} \\ \lesssim \|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\sum_{q\in\mathbb{Z}}\sum_{|q-q''|\leq 10}\|\dot{\Delta}_{q''}(\delta u,\, \nabla\delta Q)\|_{L^2_x}\|\dot{\Delta}_q(\delta u,\, \nabla\delta Q)\|_{L^2_x} \\ \lesssim \|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|(\delta u,\, \nabla\delta Q)\|_{L^2_x}^2 \\ \lesssim \|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|(\delta u,\, \nabla\delta Q)\|_{\dot{H}^{-\frac{1}{2}}}\|(\nabla\delta u,\, \Delta\delta Q)\|_{\dot{H}^{-\frac{1}{2}}} \\ \lesssim \|\nabla Q_2\|_{L^2_x}^2\|\Delta Q_2\|_{L^2_x}^2\|(\delta u,\, \nabla\delta Q)\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_\nu\|\nabla\delta u\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_{\Gamma,L}\|\Delta\delta Q\|_{\dot{H}^{-\frac{1}{2}}}^2.\end{aligned} \tag{56}$$

Now, when $j = 2$ in (54), we remark that

$$\begin{aligned}\mathcal{I}_2^1(q,q',q'') := \int_{\mathbb{R}_2}\mathrm{tr}\Big\{\Big([\dot{\Delta}_q,\, \dot{S}_{q'-1}Q_2]\mathrm{tr}\{(\dot{S}_{q''-1}Q_2 - \dot{S}_{q'-1}Q_2)\,\dot{\Delta}_{q'}\dot{\Delta}_{q''}\nabla\delta u)\}\,\mathrm{Id}\Big)\,\dot{\Delta}_q\Delta\delta Q + \\ \Big([\dot{\Delta}_q,\, \dot{S}_{q'-1}Q_2]\mathrm{tr}\{(\dot{S}_{q''-1}Q_2 - \dot{S}_{q'-1}Q_2)\,\dot{\Delta}_{q'}\dot{\Delta}_{q''}\Delta\delta Q)\}\,\mathrm{Id}\Big)\,\dot{\Delta}_q\nabla\delta u\Big\} \\ \lesssim 2^{-q}\|\dot{S}_{q'-1}\nabla Q_2\|_{L^\infty_x}\|\dot{S}_{q''-1}Q_2 - \dot{S}_{q'-1}Q_2\|_{L^\infty_x}\|\dot{\Delta}_q\dot{\Delta}_{q''}(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}\|\dot{\Delta}_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ \lesssim 2^{-q}\|\dot{S}_{q'-1}\nabla Q_2\|_{L^\infty_x}2^{-q'}\|\dot{S}_{q''-1}\nabla Q_2 - \dot{S}_{q'-1}\nabla Q_2\|_{L^\infty_x}\|\dot{\Delta}_{q''}(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}\|\dot{\Delta}_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ \lesssim 2^{-q-q'}2^{\frac{q'}{2}}\|\dot{S}_{q'-1}\nabla Q_2\|_{L^4_x}2^{\frac{q''}{2}}\|\dot{S}_{q''-1}\nabla Q_2\|_{L^4_x}2^{q''}\|\dot{\Delta}_{q''}(\delta u,\, \nabla\delta Q)\|_{L^2_x}2^q\|\dot{\Delta}_q(\delta u,\, \nabla\delta Q)\|_{L^2_x} \\ \lesssim 2^{\frac{3q''}{2}-\frac{q'}{2}}\|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|\dot{\Delta}_{q''}(\delta u,\, \nabla\delta Q)\|_{L^2_x}\|\dot{\Delta}_q(\delta u,\, \nabla\delta Q)\|_{L^2_x},\end{aligned}$$

which is equivalent to (55). Hence, proceeding as in (56), we get

$$2L\xi\sum_{q\in\mathbb{Z}}\sum_{\substack{|q-q'|\leq 5\\ |q'-q''|\leq 5}} 2^{-q}\mathcal{I}_2^1(q,q',q'') \lesssim \|\nabla Q_2\|_{L^2_x}^2\|\Delta Q_2\|_{L^2_x}^2\|(\delta u,\, \nabla\delta Q)\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_\nu\|\nabla\delta u\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_{\Gamma,L}\|\Delta\delta Q\|_{\dot{H}^{-\frac{1}{2}}}^2.$$

Concerning the term of (54) related to $j = 4$, we have

$$\begin{aligned}
\mathcal{I}_4^1(q,q',q'') := \int_{\mathbb{R}_2} \mathrm{tr}\Big\{ \Big([\dot\Delta_q,\, \dot S_{q'-1}Q_2]\mathrm{tr}\{ \dot\Delta_{q'}(\, \dot\Delta_{q''}Q_2 \dot S_{q''+2}\nabla\delta u\,)\}\,\mathrm{Id}\Big)\dot\Delta_q \Delta\delta Q - \\
- \Big([\dot\Delta_q,\, \dot S_{q'-1}Q_2]\mathrm{tr}\{ \dot\Delta_{q'}(\, \dot\Delta_{q''}Q_2 \dot S_{q''+2}\Delta\delta Q\,)\}\,\mathrm{Id}\Big)\dot\Delta_q \nabla\delta u\Big\} \\
\lesssim 2^{-q}\|\dot S_{q'-1}\nabla Q_2\|_{L^\infty_x}\|\dot\Delta_q(\, \dot\Delta_{q''}Q_2\dot S_{q''+2}\nabla\delta u,\; \dot\Delta_{q''}Q_2\dot S_{q''+2}\Delta\delta Q\,)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\
\lesssim 2^{-q}\|\dot S_{q'-1}\nabla Q_2\|_{L^\infty_x}2^q\|\dot\Delta_q(\, \dot\Delta_{q''}Q_2\dot S_{q''+2}\nabla\delta u,\; \dot\Delta_{q''}Q_2\dot S_{q''+2}\Delta\delta Q\,)\|_{L^1_x}\|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \quad (57) \\
\lesssim \|\dot S_{q'-1}\nabla Q_2\|_{L^\infty_x}\|\dot\Delta_{q''}Q_2\|_{L^2_x}\|\dot S_{q''+2}(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\
\lesssim \|\dot S_{q'-1}\nabla Q_2\|_{L^\infty_x}2^{-2q''}\|\dot\Delta_{q''}\Delta Q_2\|_{L^2_x}\|\dot S_{q''+2}(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\
\lesssim 2^{-2q''}2^{q'}\|\dot S_{q'-1}\nabla Q_2\|_{L^2_x}\|\dot\Delta_{q''}\Delta Q_2\|_{L^2_x}2^{q''}\|\dot S_{q''+2}(\delta u,\, \nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\
\lesssim 2^{q'-q''}\|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|\dot S_{q''+2}(\delta u,\, \nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}
\end{aligned}$$

Hence

$$\begin{aligned}
2L\xi\sum_{q\in\mathbb{Z}}\;\sum_{|q-q'|\le 5}\;\sum_{q''\ge q'-5} 2^{-q}\mathcal{I}_4^1(q,q',q'') \lesssim \\
\lesssim \|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\sum_{q\in\mathbb{Z}}2^{-\frac{q}{2}}\|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}\sum_{[q-q'|\le 5}2^{\frac{q'-q}{2}}\sum_{q''\ge q'-5}2^{\frac{q'-q''}{2}}2^{-\frac{q''}{2}}\|\dot S_{q''+2}(\delta u,\, \nabla\delta Q)\|_{L^2_x} \\
\lesssim \|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\sum_{q\in\mathbb{Z}}2^{-\frac{q}{2}}\|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}\sum_{q''\ge q-10}2^{\frac{q-q''}{2}}2^{-\frac{q''}{2}}\|\dot S_{q''+2}(\delta u,\, \nabla\delta Q)\|_{L^2_x} \quad (58) \\
\lesssim \|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|(\nabla\delta u,\, \Delta\delta Q)\|_{\dot H^{-\frac12}}\Big[\sum_{q\in\mathbb{Z}}\Big|\sum_{q-q''\le 10}2^{\frac{q-q''}{2}}2^{-\frac{q''+2}{2}}\|\dot S_{q''+2}(\delta u,\, \nabla\delta Q)\|_{L^2_x}\Big|^2\Big]^{\frac12} \\
\lesssim \|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|(\nabla\delta u,\, \Delta\delta Q)\|_{\dot H^{-\frac12}}\Big(\sum_{q\le 10}2^{\frac{q}{2}}\Big)\Big(\sum_{q\in\mathbb{Z}}2^{-q}\|\dot S_q(\delta u,\, \nabla\delta Q)\|_{L^2_x}^2\Big)^{\frac12} \\
\lesssim \|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|(\nabla\delta u,\, \Delta\delta Q)\|_{\dot H^{-\frac12}}\|(\delta u,\, \nabla\delta Q)\|_{\dot H^{-\frac12}} \\
\lesssim \|\nabla Q_2\|_{L^2_x}^2\|\Delta Q_2\|_{L^2_x}^2\|(\delta u,\, \nabla\delta Q)\|_{\dot H^{-\frac12}}^2 + C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2 + C_{\Gamma,L}\|\Delta\delta Q\|_{\dot H^{-\frac12}}^2.
\end{aligned}$$

Concerning (54), it remains to control the term related to $j = 3$. We fix $0 < \varepsilon \le 5/6$ and we consider the low frequencies $q \le N$, for some suitable positive $N \ge 1$ (so that $1 + \sqrt{N} < 2\sqrt{N}$):

$$\begin{aligned}
\mathcal{I}_3^1(q,q') : = \int_{\mathbb{R}_2} \mathrm{tr}\Big\{ \Big([\dot\Delta_q,\, \dot S_{q'-1}Q_2]\mathrm{tr}\{ \dot S_{q'-1}Q_2\, \dot\Delta_{q'}\nabla\delta u\}\,\mathrm{Id}\Big)\,\dot\Delta_q \Delta\delta Q - \\
- \Big([\dot\Delta_q,\, \dot S_{q'-1}Q_2]\mathrm{tr}\{ \dot S_{q'-1}Q_2\, \dot\Delta_{q'}\Delta\delta Q\}\,\mathrm{Id}\Big)\,\dot\Delta_q \nabla\delta u\Big\} \\
\lesssim 2^{-q}\|\dot S_{q'-1}\nabla Q_2\|_{L^{\frac{2}{\varepsilon}}_x}\|\dot S_{q'-1}Q_2\,\dot\Delta_{q'}\nabla\delta u,\, \dot S_{q'-1}Q_2\,\dot\Delta_{q'}\Delta\delta Q\|_{L^{\frac{2}{1-\varepsilon}}_x}\|\dot\Delta_q(\,\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\
\lesssim \|\dot S_{q'-1}\nabla Q_2\|_{L^{\frac{2}{\varepsilon}}_x}\|\dot S_{q'-1}Q_2\|_{L^\infty_x}2^{-q}\|\dot\Delta_{q'}(\,\nabla\delta u,\, \Delta\delta Q\,)\|_{L^{\frac{2}{1-\varepsilon}}_x}\|\dot\Delta_q(\,\nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}.
\end{aligned}$$

Thanks to Theorem A.2, we get

$$\|\dot S_{q'-1}Q_2\|_{L^\infty_x} \lesssim (1+\sqrt{q'-1})\|(Q_2,\, \nabla Q_2)\|_{L^2_x} \lesssim (1+\sqrt{N})\|(Q_2,\, \nabla Q_2)\|_{L^2_x} \lesssim \sqrt{N}\|(Q_2,\, \nabla Q_2)\|_{L^2_x},$$

hence $\mathcal{I}_3^1(q,q')$ is bounded by

$$\mathcal{I}_3^1(q,q') \lesssim \sqrt{N}\|(Q_2,\, \nabla Q_2)\|_{L^{\frac{2}{\varepsilon}}_x}\|(Q_2,\, \nabla Q_2)\|_{L^2_x}\|\dot\Delta_{q'}(\,\delta u,\, \nabla\delta Q\,)\|_{L^{\frac{2}{1-\varepsilon}}_x}\|\dot\Delta_q(\,\nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}. \quad (59)$$

Now, we will need the following inequality, which will finally lead to the delicate double-logarithmic estimate:

$$\|(Q_2,\, \nabla Q_2)\|_{L^{\frac{2}{\varepsilon}}_x} \le \frac{C}{\sqrt{\varepsilon}}\|(Q_2,\, \nabla Q_2)\|_{L^2_x}^{\varepsilon}\|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x}^{1-\varepsilon},$$

This is a consequence of Lemma A.3, imposing $p = 1/\varepsilon$, where C is a positive constant independent of ε and Q_2. We will see that the double-logarithmic term comes out of a suitable choice of ε in terms of N. Again, using Lemma A.3 we have

$$\begin{aligned}
\|\dot\Delta_{q'}(\,\delta u,\, \nabla\delta Q\,)\|_{L^{\frac{2}{1-\varepsilon}}_x} &\le \frac{C}{1-\varepsilon}\|\dot\Delta_{q'}(\,\delta u,\, \nabla\delta Q\,)\|_{L^2_x}^{1-\varepsilon}\|\dot\Delta_{q'}(\,\nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}^{\varepsilon} \\
&\le 6C\|\dot\Delta_{q'}(\,\delta u,\, \nabla\delta Q\,)\|_{L^2_x}^{1-\varepsilon}\|\dot\Delta_{q'}(\,\nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}^{\varepsilon},
\end{aligned}$$

since $\varepsilon \le 5/6$. Hence (59) becomes

$$\begin{aligned}\mathcal{I}_3^1(q,q') \lesssim \sqrt{\frac{N}{\varepsilon}} \|(Q_2,\, \nabla Q_2)\|_{L^2_x}^{1+\varepsilon} \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x}^{1-\varepsilon} \times \\ \times \|\dot{\Delta}_{q'}(\, \delta u,\, \nabla\delta Q\,)\|_{L^2_x}^{1-\varepsilon} \|\dot{\Delta}_{q'}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}^{\varepsilon} \|\dot{\Delta}_{q}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x},\end{aligned} \tag{60}$$

thus, since $ab \leq a^{2/(1-\varepsilon)} + b^{2/(1+\varepsilon)}$, we deduce

$$\begin{aligned}\mathcal{I}_3^1(q,q') &\lesssim \Big(\frac{N}{\varepsilon}\Big)^{\frac{1}{1-\varepsilon}} \|(Q_2,\, \nabla Q_2)\|_{L^2_x}^{\frac{2(1+\varepsilon)}{1-\varepsilon}} \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x}^2 \|\dot{\Delta}_{q'}(\, \delta u,\, \nabla\delta Q\,)\|_{L^2_x}^2 + \\ &\qquad + \min\{C_\nu,\, C_\Gamma\} \|\dot{\Delta}_{q'}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}^{\frac{2\varepsilon}{1+\varepsilon}} \|\dot{\Delta}_{q}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}^{\frac{2}{1+\varepsilon}} \\ &\lesssim \Big(\frac{N}{\varepsilon}\Big)^{\frac{1}{1-\varepsilon}} \|(Q_2,\, \nabla Q_2)\|_{L^2_x}^{\frac{2(1+\varepsilon)}{1-\varepsilon}} \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x}^2 \|\dot{\Delta}_{q'}(\, \delta u,\, \nabla\delta Q\,)\|_{L^2_x}^2 + \\ &\qquad + \min\{C_\nu,\, C_\Gamma\} \Big(\|\dot{\Delta}_{q'}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}^2 + \|\dot{\Delta}_{q}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}^2 \Big).\end{aligned}$$

Imposing $\varepsilon = (1+\ln N)^{-1}$ and observing that $\frac{1}{1-\varepsilon} = 1 + 1/\ln N$

$$N^{\frac{1}{1-\varepsilon}} = N\, N^{\frac{1}{\ln N}} = eN, \quad \varepsilon^{-\frac{1}{1-\varepsilon}} = \varepsilon^{-1}\varepsilon^{-\frac{\varepsilon}{1-\varepsilon}} = (1+\ln N) e^{\frac{\varepsilon}{1-\varepsilon}\ln\frac{1}{\varepsilon}} \lesssim (1+\ln N),$$

we obtain:

$$\begin{aligned}\mathcal{I}_3^1(q,q') \lesssim N(1+\ln N) \max\big\{ \|(Q_2,\, \nabla Q_2)\|_{L^2_x}^6, 1 \big\} \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x}^2 \|\dot{\Delta}_{q'}(\, \delta u,\, \nabla\delta Q\,)\|_{L^2_x}^2 + \\ + \min\Big\{C_\nu,\, C_{\Gamma,L}\Big\} \Big(\|\dot{\Delta}_{q'}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}^2 + \|\dot{\Delta}_{q}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x}^2 \Big),\end{aligned}$$

which yields

$$\begin{aligned}\sum_{q\leq N} \sum_{|q-q'|\leq 5} 2^{-q}\mathcal{I}_3^1(q,q') \lesssim N(1+\ln N) \max\big\{ \|(Q_2,\, \nabla Q_2)\|_{L^2_x}^6, 1 \big\} \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x}^2 \|(\, \delta u,\, \nabla\delta Q\,)\|_{\dot{H}^{-\frac{1}{2}}}^2 + \\ + C_\nu \|\nabla\delta u\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_{\Gamma,L} \|\Delta\delta Q\|_{\dot{H}^{-\frac{1}{2}}}^2.\end{aligned}$$

For the high frequencies, namely for $q > N \geq 1$, we proceed as follows:

$$\begin{aligned}\mathcal{I}_3^1(q,q') &\lesssim 2^{-q} \|\dot{S}_{q'-1}\nabla Q_2\|_{L^\infty_x} \|\dot{S}_{q'-1}Q_2\, \dot{\Delta}_{q'}\nabla\delta u,\, \dot{S}_{q'-1}Q_2\, \dot{\Delta}_{q'}\Delta\delta Q\|_{L^2_x} \|\dot{\Delta}_{q}(\, \nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ &\lesssim 2^{-q}(1+\sqrt{q'}) \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x} \|\dot{S}_{q'-1}Q_2\|_{L^\infty_x} \|\dot{\Delta}_{q'}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x} \|\dot{\Delta}_{q}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x} \\ &\lesssim 2^{-q}(1+\sqrt{q'})^2 \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x} \|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|\dot{\Delta}_{q'}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x} \|\dot{\Delta}_{q}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x} \\ &\lesssim q' \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x} \|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|\dot{\Delta}_{q'}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x} \|\dot{\Delta}_{q}(\, \delta u,\, \nabla\delta Q\,)\|_{L^2_x},\end{aligned}$$

which implies

$$\begin{aligned}&\sum_{q>N} \sum_{|q-q'|\leq 5} 2^{-q}\mathcal{I}_3^1(q,q') \\ &\qquad \lesssim \sum_{q>N} \sum_{|q-q'|\leq 5} 2^{-2q} q' \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x} \|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|\dot{\Delta}_{q'}(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{L^2_x} \|\dot{\Delta}_{q}(\, \delta u,\, \nabla\delta Q\,)\|_{L^2_x} \\ &\qquad \lesssim \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x} \|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|(\, \delta u,\, \nabla\delta Q\,)\|_{L^2_x} \|(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{\dot{H}^{-\frac{1}{2}}} \sum_{q>N} \sum_{|q-q'|\leq 5} 2^{-\frac{3}{2}q+\frac{1}{2}q'} q' \\ &\qquad \lesssim 2^{-\frac{N}{2}} \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x} \|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|(\, \delta u,\, \nabla\delta Q\,)\|_{L^2_x} \|(\, \nabla\delta u,\, \Delta\delta Q\,)\|_{\dot{H}^{-\frac{1}{2}}}.\end{aligned}$$

Summarizing, we get

$$\begin{aligned}\sum_{q\in\mathbb{Z}} \sum_{|q-q'|\leq 5} 2^{-q}\mathcal{I}_3^1(q,q') \lesssim N(1+\ln N) \max\big\{ \|(Q_2,\, \nabla Q_2)\|_{L^2_x}^6, 1 \big\} \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x}^2 \|(\, \delta u,\, \nabla\delta Q\,)\|_{\dot{H}^{-\frac{1}{2}}}^2 + \\ + 2^{-N} \|(\nabla Q_2,\, \Delta Q_2)\|_{L^2_x}^2 \|(Q_2,\, \nabla Q_2)\|_{L^2_x}^2 \|(u_1,\, u_2,\, \nabla Q_1,\, \nabla Q_2)\|_{L^2_x}^2 + C_\nu \|\nabla\delta u\|_{\dot{H}^{-\frac{1}{2}}}^2 + C_{\Gamma,L} \|\Delta\delta Q\|_{\dot{H}^{-\frac{1}{2}}}^2.\end{aligned} \tag{61}$$

Choosing $N = N(t) := \lceil \ln(1+e+1/\Phi(t)) \rceil$ (thus $\varepsilon < 1/(1+\ln\ln\{1+e\}) < 5/6$) where with $\lceil\cdot\rceil$ we denote the ceiling function, relation (61) implies

$$\begin{aligned}
&\sum_{q\in\mathbb{Z}}\sum_{|q-q'|\le 5} 2^{-q}\mathcal{I}_1^3(q,q')\\
&\lesssim \max\big\{\|(Q_2,\,\nabla Q_2)\|_{L^2_x}^6, 1\big\}\|(\nabla Q_2,\,\Delta Q_2)\|_{L^2_x}^2\|(\,\delta u,\,\nabla\delta Q\,)\|_{\dot H^{-\frac12}}^2 \ln\Big(e+\frac{1}{\Phi(t)}\Big)\Big(1+\ln\ln\Big(e+\frac{1}{\Phi(t)}\Big)\Big)+ \qquad (62)\\
&+\|(\nabla Q_2,\,\Delta Q_2)\|_{L^2_x}^2\|(Q_2,\,\nabla Q_2)\|_{L^2_x}^2\|(u_1,\,u_2,\,\nabla Q_1,\,\nabla Q_2)\|_{L^2_x}^2\Phi(t)+C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2+C_{\Gamma,L}\|\Delta\delta Q\|_{\dot H^{-\frac12}}^2.
\end{aligned}$$

Estimate of $\mathcal{J}_q^2$ Now, we handle the term of (53) related to $i=2$, namely

$$\begin{aligned}
\sum_{j=1}^4\sum_{|q-q'|\le 5}\int_{\mathbb{R}_2}\mathrm{tr}\{\,(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\mathrm{tr}\{\dot\Delta_q\mathcal{J}^j_{q'}(Q_2,\,\nabla\delta u)\}\dot\Delta_q\Delta\delta Q+\\
(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\mathrm{tr}\{\dot\Delta_q\mathcal{J}^j_{q'}(Q_2,\,\Delta\delta Q)\}\dot\Delta_q\nabla\delta u\,\}.
\end{aligned}\tag{63}$$

When $j=1$, we have

$$\begin{aligned}
\mathcal{I}_1^2(q,q',q''):=\int_{\mathbb{R}_2}&\mathrm{tr}\Big\{(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\mathrm{tr}\{\dot\Delta_q\big(\,[\dot\Delta_{q'},\dot S_{q''-1}Q_2]\dot\Delta_{q''}\nabla\delta u\,\big)\}\dot\Delta_q\Delta\delta Q+\\
&-(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\mathrm{tr}\{\dot\Delta_q\big(\,[\dot\Delta_{q'},\dot S_{q''-1}Q_2]\dot\Delta_{q''}\Delta\delta Q\,\big)\}\dot\Delta_q\nabla\delta u\Big\}\\
&\lesssim\|\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot\Delta_q\big(\,[\dot\Delta_{q'},\dot S_{q''-1}Q_2]\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\,\big)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^{-\frac q2}\|\dot S_{q'-1}\nabla Q_2-\dot S_{q-1}\nabla Q_2\|_{L^4_x}2^{\frac q2}\|\dot\Delta_q\big(\,[\dot\Delta_{q'},\dot S_{q''-1}Q_2]\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\,\big)\|_{L^{\frac43}_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim\|\nabla Q_2\|_{L^4_x}2^{-q'}\|\dot S_{q''-1}\nabla Q_2\|_{L^4_x}\|\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^{-q'+q''+q}\|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|\dot\Delta_{q''}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\delta u,\,\nabla\delta Q)\|_{L^2_x}.
\end{aligned}$$

Since $|q-q'|\le 5$ and $|q'-q''|\le 5$ then $-q'+q''+q\simeq 3q''/2-q'/2$, so that the last inequality is equivalent to (55). Hence, proceeding as in (56), we get

$$2L\xi\sum_{q\in\mathbb{Z}}\sum_{\substack{|q-q'|\le 5\\|q'-q''|\le 5}}2^{-q}\mathcal{I}_1^2(q,q',q'')\lesssim\|\nabla Q_2\|_{L^2_x}^2\|\Delta Q_2\|_{L^2_x}^2\|(\delta u,\,\nabla\delta Q)\|_{\dot H^{-\frac12}}^2+C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2+C_{\Gamma,L}\|\Delta\delta Q\|_{\dot H^{-\frac12}}^2.$$

When $j=2$, we observe that

$$\begin{aligned}
\mathcal{I}_2^2(q,q',q''):=\int_{\mathbb{R}_2}&\mathrm{tr}\Big\{(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\mathrm{tr}\{\,(\dot S_{q''-1}Q_2-\dot S_{q'-1}Q_2)\dot\Delta_{q'}\dot\Delta_{q''}\nabla\delta u\,\}\dot\Delta_q\Delta\delta Q+\\
&-(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\mathrm{tr}\{(\dot S_{q''-1}Q_2-\dot S_{q'-1}Q_2)\dot\Delta_{q'}\dot\Delta_{q''}\Delta\delta Q\,\}\dot\Delta_q\nabla\delta u\Big\}\\
\lesssim\ &\|\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot S_{q''-1}Q_2-\dot S_{q'-1}Q_2\|_{L^\infty_x}\|\dot\Delta_{q'}\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
\lesssim\ &2^{-\frac{q+q'}{2}}\|\dot S_{q'-1}\nabla Q_2-\dot S_{q-1}\nabla Q_2\|_{L^4_x}\|\dot S_{q''-1}\nabla Q_2-\dot S_{q'-1}\nabla Q_2\|_{L^4_x}\|\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
\lesssim\ &2^{\frac{q'}{2}+\frac q2}\|\nabla Q_2\|_{L^4_x}^2\|\dot\Delta_{q''}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
\lesssim\ &2^{\frac{q'}{2}+\frac q2}\|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|\dot\Delta_{q'}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\delta u,\,\nabla\delta Q)\|_{L^2_x}.
\end{aligned}$$

Since $|q-q'|\le 5$ and $|q'-q''|\le 5$ then $q'/2+q/2\simeq 3q''/2-q'/2$, so that the last inequality is equivalent to (55). Hence, proceeding as in (56), we get

$$2L\xi\sum_{q\in\mathbb{Z}}\sum_{\substack{|q-q'|\le 5\\|q'-q''|\le 5}}2^{-q}\mathcal{I}_2^2(q,q',q'')\lesssim\|\nabla Q_2\|_{L^2_x}^2\|\Delta Q_2\|_{L^2_x}^2\|(\delta u,\,\nabla\delta Q)\|_{\dot H^{-\frac12}}^2+C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2+C_{\Gamma,L}\|\Delta\delta Q\|_{\dot H^{-\frac12}}^2.$$

When $j=4$:

$$\begin{aligned}
\mathcal{I}_4^2(q,q',q''):=\int_{\mathbb{R}_2}&\Big\{(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\mathrm{tr}\{\,\dot\Delta_{q'}(\,\dot\Delta_{q''}Q_2\dot S_{q''+2}\nabla\delta u\,)\,\}\dot\Delta_q\Delta\delta Q+\\
&-(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\mathrm{tr}\{\,\dot\Delta_{q'}(\,\dot\Delta_{q''}Q_2\dot S_{q''+2}\Delta\delta Q\,)\,\}\dot\Delta_q\nabla\delta u\Big\}\\
&\lesssim\|\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot\Delta_{q'}(\,\dot\Delta_{q''}Q_2\dot S_{q''+2}(\nabla\delta u,\,\Delta\delta Q\,)\,)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^{q'}\|\dot S_{q'-1}\nabla Q_2-\dot S_{q-1}\nabla Q_2\|_{L^2_x}\|\dot\Delta_{q'}(\,\dot\Delta_{q''}Q_2\dot S_{q''+2}(\nabla\delta u,\,\Delta\delta Q\,)\,)\|_{L^1_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}
\end{aligned}$$

$$
\begin{aligned}
&\lesssim 2^{q'-q''}\|\nabla Q_2\|_{L^2_x}\|\dot\Delta_{q''}\Delta Q_2\|_{L^2_x}\|\dot S_{q''+2}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^{q'-q''}\|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|\dot S_{q''+2}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x},
\end{aligned}
$$

which is equivalent to the last inequality of (57). Thus, arguing as in (58), we deduce

$$
2L\xi\sum_{q\in\mathbb{Z}}\sum_{\substack{|q-q'|\le 5\\ q''\ge q'-5}}2^{-q}\mathcal{I}_4^2(q,q',q'')\lesssim\|\nabla Q_2\|_{L^2_x}^2\|\Delta Q_2\|_{L^2_x}^2\|(\delta u,\,\nabla\delta Q)\|^2_{\dot H^{-\frac12}}+C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}.
$$

When $j=3$ we fix a real number $N>1$ and we consider the low frequencies $q'\le N$ as follows

$$
\begin{aligned}
\mathcal{I}_3^2(q,q') := \int_{\mathbb{R}_2}\operatorname{tr}\Big\{&(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\operatorname{tr}\{\,\dot\Delta_q(\,\dot S_{q'-1}Q_2\,\dot\Delta_{q'}\nabla\delta u\,)\,\}\dot\Delta_q\Delta\delta Q+\\
&-(\,\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\,)\operatorname{tr}\{\,\dot\Delta_q(\,\dot S_{q'-1}Q_2\,\dot\Delta_{q'}\Delta\delta Q\,)\}\dot\Delta_q\nabla\delta u\Big\}\\
\lesssim{}&\|\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot\Delta_q(\,\dot S_{q'-1}Q_2\,\dot\Delta_{q'}(\nabla\delta u,\,\Delta\delta Q)\,)\,\}\|_{L^2_x}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
\lesssim{}&2^{-q}\|\dot S_{q'-1}\Delta Q_2-\dot S_{q-1}\Delta Q_2\|_{L^2_x}\|\dot S_{q'-1}Q_2\|_{L^\infty_x}\|\dot\Delta_{q'}(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\\
\lesssim{}&\|\dot S_{q'-1}\Delta Q_2-\dot S_{q-1}\Delta Q_2\|_{L^2_x}\|\dot S_{q'-1}Q_2\|_{L^\infty_x}\|\dot\Delta_{q'}(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\|\dot\Delta_q(\,\delta u,\,\nabla\delta Q\,)\|_{L^2_x}.
\end{aligned}
\tag{64}
$$

If $q'\le 1$ then $\|\dot S_{q'-1}Q_2\|_{L^\infty_x}\lesssim 2^{\frac{q'}{2}}\|\dot S_{q'-1}Q_2\|_{L^2_x}\le\|Q_2\|_{L^2_x}$, while if $1<q'\le N$ we have

$$
\|\dot S_{q'-1}Q_2\|_{L^\infty_x}\lesssim(\|Q_2\|_{L^2_x}+\sqrt{q'-1}\|\nabla Q_2\|_{L^2_x})\lesssim(\|Q_2\|_{L^2_x}+\sqrt{N}\|\nabla Q_2\|_{L^2_x}),
$$

thanks to Theorem A.2. Therefore, we deduce that

$$
\begin{aligned}
\mathcal{I}_3^2(q,q')&\lesssim\|\Delta Q_2\|_{L^2_x}(\|Q_2\|_{L^2_x}+\sqrt{N}\|\nabla Q_2\|_{L^2_x})\|\dot\Delta_{q'}(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\|\dot\Delta_q(\,\delta u,\,\nabla\delta Q\,)\|_{L^2_x}\\
&\lesssim(1+N)\|\Delta Q_2\|^2_{L^2_x}\|(Q_2,\,\nabla Q_2)\|^2_{L^2_x}\|\dot\Delta_q(\,\delta u,\,\nabla\delta Q\,)\|^2_{L^2_x}+C_\nu\|\dot\Delta_{q'}\nabla\delta u\|^2_{L^2_x}+C_{\Gamma,L}\|\dot\Delta_{q'}\Delta\delta Q\|^2_{L^2_x}.
\end{aligned}
$$

Hence

$$
\begin{aligned}
\sum_{q'\le N}\sum_{|q'-q|\le 5}2^{-q}\mathcal{I}_3^2(q,q')\lesssim{}&(1+N)\|\Delta Q_2\|^2_{L^2_x}\|(Q_2,\,\nabla Q_2)\|^2_{L^2_x}\|(\,\delta u,\,\nabla\delta Q\,)\|^2_{\dot H^{-\frac12}}+\\
&+C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}.
\end{aligned}
\tag{65}
$$

For the high frequencies $q'>N$ we get,

$$
\begin{aligned}
\mathcal{I}_3^2(q,q')&\lesssim\|\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot\Delta_q(\,\dot S_{q'-1}Q_2\,\dot\Delta_{q'}(\nabla\delta u,\,\Delta\delta Q)\,)\,\}\|_{L^2_x}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^{-q}\|\dot S_{q'-1}\Delta Q_2-\dot S_{q-1}\Delta Q_2\|_{L^2_x}\|\dot S_{q'-1}Q_2\|_{L^\infty_x}\|\dot\Delta_{q'}(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\\
&\lesssim 2^{\frac{q'-q}{2}}\|\Delta Q_2\|_{L^2_x}(1+\sqrt{q'-1})\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|\dot\Delta_{q'}(\,\delta u,\,\nabla\delta Q\,)\|_{L^2_x}\|(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{\dot H^{-\frac12}}\\
&\lesssim(1+\sqrt{q'-1})\|\Delta Q_2\|_{L^2_x}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|(\,\delta u,\;\;\delta Q\,)\|_{L^2_x}\|(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{\dot H^{-\frac12}},
\end{aligned}
\tag{66}
$$

therefore

$$
\begin{aligned}
\sum_{q'>N}\sum_{|q-q'|\le 5}2^{-q}\mathcal{I}_3^2(q,q')&\lesssim 2^{-N}\|\Delta Q_2\|_{L^2_x}\|\nabla Q_2\|_{L^2_x}\|(\,\delta u,\;\;\delta Q\,)\|_{L^2_x}\|(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{\dot H^{-\frac12}}\\
&\lesssim 2^{-2N}\|\Delta Q_2\|^2_{L^2_x}\|\nabla Q_2\|^2_{L^2_x}\|(\,\delta u,\;\;\delta Q\,)\|^2_{L^2_x}+C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}.
\end{aligned}
\tag{67}
$$

Summarizing (65) and (67), we get

$$
\begin{aligned}
\sum_{q'\in\mathbb{Z}}\sum_{|q'-q|\le 5}2^{-q}\mathcal{I}_3^2(q,q')\lesssim{}&(1+N)\|\Delta Q_2\|^2_{L^2_x}\|(Q_2,\,\nabla Q_2)\|^2_{L^2_x}\|(\,\delta u,\,\nabla\delta Q\,)\|^2_{\dot H^{-\frac12}}+\\
&+2^{-2N}\|\Delta Q_2\|^2_{L^2_x}\|\nabla Q_2\|^2_{L^2_x}\|(\,\delta u,\;\;\delta Q\,)\|^2_{L^2_x}+C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}.
\end{aligned}
\tag{68}
$$

Now we define $N:=\lceil\ln\{e+1/\Phi(t)\}/2\rceil$, obtaining

$$
\begin{aligned}
\sum_{q\in\mathbb{Z}}\sum_{|q'-q|\le 5}2^{-q}\mathcal{I}_3^2(q,q')\lesssim{}&\|\Delta Q_2(t)\|^2_{L^2_x}\|(Q_2,\,\nabla Q_2)(t)\|^2_{L^2_x}\|(\,\delta u,\,\nabla\delta Q(t)\,)\|^2_{\dot H^{-\frac12}}+C_\nu\|\nabla\delta u(t)\|^2_{\dot H^{-\frac12}}+\\
&+C_{\Gamma,L}\|\Delta\delta Q(t)\|^2_{\dot H^{-\frac12}}+\|\Delta Q_2(t)\|^2_{L^2_x}\|\nabla Q_2(t)\|^2_{L^2_x}\|(\,\delta u,\;\;\delta Q\,)(t)\|^2_{L^2_x}\Big(1+\ln\Big(e+\frac{1}{\Phi(t)}\Big)\Big).
\end{aligned}
\tag{69}
$$

Estimate of $\mathcal{J}_q^3$ Now, let us deal with the term of (53) related to $i=3$, namely

$$\begin{aligned}&\int_{\mathbb{R}_2}\mathrm{tr}\{\dot S_{q-1}Q_2\mathrm{tr}\{\dot\Delta_q(Q_2\nabla\delta u)\}\dot\Delta_q\Delta\delta Q-\dot S_{q-1}Q_2]\mathrm{tr}\{\dot\Delta_q(Q_2\Delta\delta Q)\}\dot\Delta_q\nabla\delta u\,\}=\\&\qquad=\sum_{j=1}^{4}\int_{\mathbb{R}_2}\mathrm{tr}\{\,\dot S_{q-1}Q_2\mathrm{tr}\{\mathcal{J}^j_{q'}(Q_2,\,\nabla\delta u)\}\dot\Delta_q\Delta\delta Q-\dot S_{q-1}Q_2\mathrm{tr}\{\mathcal{J}^j_{q'}(Q_2,\,\Delta\delta Q)\}\dot\Delta_q\nabla\delta u\,\}.\end{aligned}\tag{70}$$

Let us consider $j=1$ and define

$$\mathcal{I}^3_1(q,q'):=\int_{\mathbb{R}_2}\mathrm{tr}\Big\{\dot S_{q-1}Q_2\mathrm{tr}\{[\dot\Delta_q,\dot S_{q'-1}Q_2]\dot\Delta_{q'}\nabla\delta u)\}\dot\Delta_q\Delta\delta Q-\dot S_{q-1}Q_2\mathrm{tr}\{[\dot\Delta_q,\dot S_{q'-1}Q_2]\dot\Delta_{q'}\Delta\delta Q)\}\dot\Delta_q\nabla\delta u\Big\}.$$

We proceed as for proving (60): we fix a positive real $\varepsilon\in(0,5/6]$ and we consider the low frequencies $q\le N$, for a suitable positive $N\ge 1$.

$$\begin{aligned}\mathcal{I}^3_1(q,q')&=\int_{\mathbb{R}_2}\mathrm{tr}\Big\{\dot S_{q-1}Q_2\mathrm{tr}\{[\dot\Delta_q,\,\dot S_{q'-1}Q_2]\,\dot\Delta_{q'}\nabla\delta u\}\dot\Delta_q\Delta\delta Q-\dot S_{q-1}Q_2\mathrm{tr}\{[\dot\Delta_q,\,\dot S_{q'-1}Q_2]\,\dot\Delta_{q'}\Delta\delta Q\}\dot\Delta_q\nabla\delta u\Big\}\\&\lesssim 2^{-q'}\|\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot S_{q'-1}\nabla Q_2\|_{L^{\frac{2}{\varepsilon}}_x}\|\dot\Delta_{q'}(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L_x^{\frac{2}{1-\varepsilon}}}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\\&\lesssim(1+\sqrt{N})2^{q-q'}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|\dot S_{q'-1}\nabla Q_2\|_{L^{\frac{2}{\varepsilon}}_x}\|\dot\Delta_{q'}(\,\delta u,\,\nabla\delta Q\,)\|_{L_x^{\frac{2}{1-\varepsilon}}}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\\&\lesssim\sqrt{\frac{N}{\varepsilon}}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|\dot S_{q'-1}\nabla Q_2\|^{\varepsilon}_{L^2_x}\|\dot S_{q'-1}\Delta Q_2\|^{1-\varepsilon}_{L^2_x}\times\\&\qquad\qquad\times\|\dot\Delta_{q'}(\,\delta u,\,\nabla\delta Q\,)\|^{1-\varepsilon}_{L^2_x}\|\dot\Delta_{q'}(\,\nabla\delta u,\,\Delta\delta Q\,)\|^{\varepsilon}_{L^2_x}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x},\end{aligned}$$

which is equivalent to the last inequality of (60). Hence, arguing as for proving (62), we get

$$\begin{aligned}&\sum_{q\in\mathbb{Z}}\sum_{|q-q'|\le5}2^{-q}\mathcal{I}^3_1(q,q')\\&\lesssim\max\big\{\|(Q_2,\,\nabla Q_2)\|^6_{L^2_x},1\big\}\|(\nabla Q_2,\,\Delta Q_2)\|^2_{L^2_x}\|(\,\delta u,\,\nabla\delta Q\,)\|^2_{\dot H^{-\frac12}}\ln\Big(1+e+\frac{1}{\Phi(t)}\Big)\Big(1+\ln\ln\Big(1+e+\frac{1}{\Phi(t)}\Big)\Big)+\\&+\|(\nabla Q_2,\,\Delta Q_2)\|^2_{L^2_x}\|(Q_2,\,\nabla Q_2)\|^2_{L^2_x}\|(u_1,\,u_2,\,\nabla Q_1,\,\nabla Q_2)\|^2_{L^2_x}\Phi(t)+C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}},\end{aligned}$$

Further on, when $j=2$ in (70), let us consider the low frequencies $q\le N$:

$$\begin{aligned}\mathcal{I}^3_2(q,q')&:=\int_{\mathbb{R}_2}\mathrm{tr}\Big\{\dot S_{q-1}Q_2\mathrm{tr}\{(\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2)\,\dot\Delta_q\dot\Delta_{q'}\nabla\delta u)\}\dot\Delta_q\Delta\delta Q+\\&\qquad\qquad-\dot S_{q-1}Q_2\mathrm{tr}\{(\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2)\,\dot\Delta_{q'}\dot\Delta_{q'}\Delta\delta Q)\}\dot\Delta_q\nabla\delta u\Big\}\\&\lesssim\|\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot\Delta_q\dot\Delta_{q'}(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\\&\lesssim\|\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot S_{q'-1}\Delta Q_2-\dot S_{q-1}\Delta Q_2\|_{L^2_x}\|\dot\Delta_{q'}(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\|\dot\Delta_q(\,\delta u,\,\nabla\delta Q\,)\|_{L^2_x},\end{aligned}$$

which is as the last inequalities of (64) (recalling that $q\sim q'$). Moreover for the high frequencies $q>N$

$$\begin{aligned}\mathcal{I}^3_2(q,q')&\lesssim\|\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot S_{q'-1}Q_2-\dot S_{q-1}Q_2\|_{L^\infty_x}\|\dot\Delta_q\dot\Delta_{q'}(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\\&\lesssim(1+\sqrt{q-1})\|(Q_2,\,\nabla Q_2)\|_{L^2_x}2^{-q}\|\dot S_{q'-1}\Delta Q_2-\dot S_{q-1}\Delta Q_2\|_{L^2_x}\times\\&\qquad\qquad\times\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\|\dot\Delta_q(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{L^2_x}\\&\lesssim(1+\sqrt{q-1})\|\Delta Q_2\|_{L^2_x}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|(\,\delta u,\,\nabla\delta Q\,)\|_{L^2_x}\|(\,\nabla\delta u,\,\Delta\delta Q\,)\|_{\dot H^{-\frac12}},\end{aligned}$$

which is the equivalent to the last inequality (66). Hence, arguing as for proving (69), we get

$$\begin{aligned}\sum_{q\in\mathbb{Z}}\sum_{|q'-q|\le5}2^{-q}\mathcal{I}^3_2(q,q')&\lesssim\|\Delta Q_2(t)\|^2_{L^2_x}\|(Q_2,\,\nabla Q_2)(t)\|^2_{L^2_x}\|(\,\delta u,\,\nabla\delta Q(t)\,)\|^2_{\dot H^{-\frac12}}+C_\nu\|\nabla\delta u(t)\|^2_{\dot H^{-\frac12}}+\\&+C_{\Gamma,L}\|\Delta\delta Q(t)\|^2_{\dot H^{-\frac12}}+\|\Delta Q_2(t)\|^2_{L^2_x}\|\nabla Q_2(t)\|^2_{L^2_x}\|(\,\delta u,\;\delta Q\,)(t)\|^2_{L^2_x}\Phi(t)\Big(1+\ln\Big(1+e+\frac{1}{\Phi(t)}\Big)\Big).\end{aligned}$$

Now, when $j=3$ in (70), we observe that

$$\mathcal{I}^3_3(q):=\int_{\mathbb{R}_2}\Big\{\mathrm{tr}\{\,\dot S_{q-1}Q_2\dot\Delta_q\nabla\delta u\,\}\mathrm{tr}\{\,\dot S_{q-1}Q_2\dot\Delta_q\Delta\delta Q\,\}-\mathrm{tr}\{\,\dot S_{q-1}Q_2\dot\Delta_q\Delta\delta Q\,\}\mathrm{tr}\{\,\dot S_{q-1}Q_2\dot\Delta_q\nabla\delta u\,\}\Big\}=0,$$

for any $q\in\mathbb{Z}$.

Thus it remains to control the $j = 4$ term, namely

$$\begin{aligned}\mathcal{I}_4^3(q,q') &:= \int_{\mathbb{R}_2} \Big\{ \dot S_{q-1} Q_2 \mathrm{tr}\{(\dot\Delta_q(\,\dot\Delta_{q'} Q_2 \dot S_{q'+2} \nabla\delta u\,)\}\dot\Delta_q \Delta\delta Q - \dot S_{q-1} Q_2 \mathrm{tr}\{(\dot\Delta_q(\,\dot\Delta_{q'} Q_2 \dot S_{q'+2}\Delta\delta Q\,)\}\dot\Delta_q \nabla\delta u \Big\} \\ &\lesssim \|\dot S_{q-1} Q_2\|_{L^\infty_x} \|\dot\Delta_q(\,\dot\Delta_{q'} Q_2 \dot S_{q'+2}(\nabla\delta u,\, \Delta\delta Q)\,)\|_{L^2_x} \|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}.\end{aligned}$$

At first let us consider the low frequencies $q \le N$, with $N > 1$:

$$\begin{aligned}\mathcal{I}_4^3(q,q') &\lesssim (1+\sqrt{N})\|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|\dot\Delta_{q'} Q_2\|_{L^\infty_x} \|\dot S_{q'+2}(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ &\lesssim (1+\sqrt{N})\|(Q_2,\, \nabla Q_2)\|_{L^2_x} 2^{-q'} \|\dot\Delta_{q'} \Delta Q_2\|_{L^2_x} \|\dot S_{q'+2}(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ &\lesssim (1+\sqrt{N})\|\nabla Q_2\|_{L^2_x} \|\dot\Delta_{q'} \Delta Q_2\|_{L^2_x} \|\dot S_{q'+2}(\delta u,\, \nabla\delta Q)\|_{L^2_x} \|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ &\lesssim (1+\sqrt{N}) 2^{\frac{q'}{2}} \|\nabla Q_2\|_{L^2_x} \|\dot\Delta_{q'} \Delta Q_2\|_{L^2_x} \|(\delta u,\, \nabla\delta Q)\|_{\dot H^{-\frac12}} \|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x},\end{aligned}$$

which yields

$$\begin{aligned}&\sum_{q\le N}\sum_{q'\ge q-5} 2^{-q}\mathcal{I}_4^3(q,q') \\ &\quad\lesssim (1+\sqrt{N})\|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|(\delta u,\, \nabla\delta Q)\|_{\dot H^{-\frac12}} \sum_{q\le N}\sum_{q'\ge q-5} 2^{\frac{q'}{2}-q} \|\dot\Delta_{q'} \Delta Q_2\|_{L^2_x} \|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ &\quad\lesssim (1+\sqrt{N})\|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|(\delta u,\, \nabla\delta Q)\|_{\dot H^{-\frac12}} \sum_{q\in\mathbb{Z}} 2^{-\frac{q}{2}} \|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \sum_{q'\ge q-5} 2^{\frac{q'-q}{2}} \|\dot\Delta_{q'} \Delta Q_2\|_{L^2_x} \\ &\quad\lesssim (1+\sqrt{N})\|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|(\delta u,\, \nabla\delta Q)\|_{\dot H^{-\frac12}} \times \\ &\qquad\qquad\times \Big(\sum_{q'\in\mathbb{Z}} \Big| \sum_{q\in\mathbb{Z}} 2^{\frac{q-q'}{2}} 1_{(-\infty,5]}(q-q') \|\dot\Delta_q \Delta Q_2\|_{L^2_x} \Big|^2 \Big)^{\frac12} \|(\nabla\delta u,\, \Delta\delta Q)\|_{\dot H^{-\frac12}},\end{aligned}$$

thus by convolution

$$\begin{aligned}&\sum_{q\le N}\sum_{q'\ge q-5} 2^{-q}\mathcal{I}_4^3(q,q')(1+\sqrt{N})\|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|\Delta Q_2\|_{L^2_x} \|(\delta u,\, \nabla\delta Q)\|_{\dot H^{-\frac12}} \|(\nabla\delta u,\, \Delta\delta Q)\|_{\dot H^{-\frac12}} \\ &\quad\lesssim (1+N)\|(Q_2,\, \nabla Q_2)\|_{L^2_x}^2 \|\Delta Q_2\|_{L^2_x}^2 \|(\delta u,\, \nabla\delta Q)\|_{\dot H^{-\frac12}}^2 + C_\nu \|\nabla\delta u\|_{\dot H^{-\frac12}}^2 + C_{\Gamma,L} \|\Delta\delta Q\|_{\dot H^{-\frac12}}^2.\end{aligned}$$

For the high frequencies, $q > N$,

$$\begin{aligned}&\sum_{q\ge N}\sum_{q'\ge q-5} 2^{-q}\mathcal{I}_2^3(q,q') \lesssim \\ &\lesssim \sum_{q\ge N}\sum_{q'\ge q-5} 2^{-q}(1+\sqrt{q-1})\|\dot S_{q-1}(Q_2,\, \nabla Q_2)\|_{L^2_x} \|\dot\Delta_{q'} Q_2\|_{L^\infty_x} \|\dot S_{q'+2}(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \|\dot\Delta_q(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ &\lesssim \|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \sum_{q\ge N} 2^{-\frac{q}{2}}(1+\sqrt{q}) \sum_{q'\ge q-5} 2^{\frac{q'-q}{2}} \|\dot\Delta_{q'} \nabla Q_2\|_{L^2_x} 2^{-\frac{q'}{2}} \|\dot S_{q'+2}(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \\ &\lesssim \|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \|(\nabla\delta u,\, \Delta\delta Q)\|_{\dot H^{-\frac12}} 2^{-\frac{N}{2}} \Big(\sum_{q\in\mathbb{Z}} \Big| \sum_{q'\ge q-5} 2^{\frac{q'-q}{2}} \|\dot\Delta_{q'} \nabla Q_2\|_{L^2_x} \Big|^2 \Big)^{\frac12},\end{aligned}$$

so that, by convolution

$$\sum_{q\ge N}\sum_{q'\ge q-5} 2^{-q}\mathcal{I}_2^3(q,q') \lesssim 2^{-N} \|(Q_2,\, \nabla Q_2)\|_{L^2_x} \|\nabla Q_2\|_{L^2_x} \|(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x} \|(\nabla\delta u,\, \Delta\delta Q)\|_{\dot H^{-\frac12}}.$$

Summarizing, we get

$$\begin{aligned}\sum_{q\in\mathbb{Z}}\sum_{q'\ge q-5} 2^{-q}\mathcal{I}_2^3(q,q') &\lesssim (1+N)\|(Q_2,\, \nabla Q_2)\|_{L^2_x}^2 \|\Delta Q_2\|_{L^2_x}^2 \|(\delta u,\, \nabla\delta Q)\|_{\dot H^{-\frac12}}^2 + C_\nu \|\nabla\delta u\|_{\dot H^{-\frac12}}^2 + \\ &\quad + C_{\Gamma,L} \|\Delta\delta Q\|_{\dot H^{-\frac12}}^2 + 2^{-2N} \|(Q_2,\, \nabla Q_2)\|_{L^2_x}^2 \|\nabla Q_2\|_{L^2_x}^2 \|(\nabla\delta u,\, \Delta\delta Q)\|_{L^2_x}^2,\end{aligned}$$

which is similar to (68), hence we can conclude as in (69).

Estimate of $\mathcal{J}^4_q$ Now, we handle the last term of (53), which is related to $i = 4$, namely

$$\begin{aligned}
&\sum_{q'\geq q-5}\int_{\mathbb{R}_2}\mathrm{tr}\big\{\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot S_{q'+2}(Q_2\nabla\delta u)\}\,\big]\dot\Delta_q\Delta\delta Q - \dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot S_{q'+2}(Q_2\Delta\delta Q)\}\,\big]\dot\Delta_q\nabla\delta u\big\} = \\
&=\sum_{q'\leq q-5}\;\sum_{q''\leq q'+1}\int_{\mathbb{R}_2}\mathrm{tr}\big\{\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot\Delta_{q'}(Q_2\nabla\delta u)\}\,\big]\dot\Delta_q\Delta\delta Q - \dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot\Delta_{q'}(Q_2\Delta\delta Q)\}\,\big]\dot\Delta_q\nabla\delta u\big\} \\
&=\sum_{j=1}^{4}\sum_{q'\leq q-5}\;\sum_{q''\leq q'+1}\int_{\mathbb{R}_2}\mathrm{tr}\big\{\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\mathcal{J}^j_{q''}(Q_2,\,\nabla\delta u)\}\,\big]\dot\Delta_q\Delta\delta Q - \\
&\qquad\qquad - \dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\mathcal{J}^j_{q''}(Q_2,\,\Delta\delta Q)\}\,\big]\dot\Delta_q\nabla\delta u\big\}.
\end{aligned}\tag{71}$$

First, we consider the term related to $j = 1$, that is

$$\begin{aligned}
\mathcal{I}^4_1(q,q',q'',q''') : &= \int_{\mathbb{R}_2}\mathrm{tr}\Big\{\;\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{[\dot\Delta_{q''},\,\dot S_{q'''-1}Q_2]\,\dot\Delta_{q'''}\nabla\delta u\}\,\big]\dot\Delta_q\Delta\delta Q + \\
&\qquad - \dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{[\dot\Delta_{q''},\,\dot S_{q'''-1}Q_2]\,\dot\Delta_{q''}\Delta\delta Q\}\,\big]\dot\Delta_q\nabla\delta u\;\Big\} \\
&\lesssim \|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{[\dot\Delta_{q''},\,S_{q'''-1}Q_2]\,\dot\Delta_{q'''}(\nabla\delta u,\,\Delta\delta Q)\,\}\,\big]\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x} \\
&\lesssim 2^q\|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{[\dot\Delta_{q''},\,S_{q'''-1}Q_2]\,\dot\Delta_{q'''}(\nabla\delta u,\,\Delta\delta Q)\,\}\,\big]\|_{L^1_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x} \\
&\lesssim 2^{q-q''}\|\dot\Delta_{q'}Q_2\|_{L^\infty_x}\|\dot S_{q'''-1}\nabla Q_2\|_{L^2_x}\|\dot\Delta_{q'''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x} \\
&\lesssim 2^{q-q'-q''}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\|\dot S_{q'''-1}\nabla Q_2\|_{L^2_x}\|\dot\Delta_{q'''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x} \\
&\lesssim 2^{q-q'-q''+q'''}\|\Delta Q_2\|_{L^2_x}\|\nabla Q_2\|_{L^2_x}\|\dot\Delta_{q'''}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}.
\end{aligned}\tag{72}$$

Hence, taking the sum in q, q', q'' and q''' (and observing that $|q''-q'''|\leq 5$), we get

$$\begin{aligned}
&\sum_{q\in\mathbb{Z}}\;\sum_{q'\geq q-5}\;\sum_{q''\leq q'+1}\;\sum_{|q'''-q''|\leq 5} 2^{-q}\mathcal{I}^4_1(q,q',q'',q''') \lesssim \\
&\lesssim \|\nabla Q_2\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\sum_{q,\,q',\,q''} 2^{\frac{q}{2}-q'}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\|\dot\Delta_{q''}(\delta u,\,\nabla\delta Q)\|_{L^2_x} \\
&\lesssim \|\nabla Q_2\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\sum_{q\in\mathbb{Z}}2^{\frac{q}{2}}\sum_{q'\geq q-5}2^{-q'}\sum_{q''\leq q'+1}2^{\frac{q''}{2}}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}2^{-\frac{q''}{2}}\|\dot\Delta_{q''}(\delta u,\,\nabla\delta Q)\|_{L^2_x} \\
&\lesssim \|\nabla Q_2\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\sum_{q''\in\mathbb{Z}}2^{-\frac{q''}{2}}\|\dot\Delta_{q''}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\sum_{q'\geq q''+1}2^{\frac{q''}{2}-q'}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\sum_{q\leq q'+5}2^{\frac{q}{2}} \\
&\lesssim \|\nabla Q_2\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\sum_{q''\in\mathbb{Z}}2^{-\frac{q''}{2}}\|\dot\Delta_{q''}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\sum_{q'\geq q''+1}2^{\frac{q''}{2}-\frac{q'}{2}}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x} \\
&\lesssim \|\nabla Q_2\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|(\delta u,\,\nabla\delta Q)\|_{\dot H^{-\frac12}}\Big(\sum_{q''\in\mathbb{Z}}\Big|\sum_{q'\geq q''+1}2^{\frac{q''}{2}-\frac{q'}{2}}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\Big|^2\Big)^{\frac12} \\
&\lesssim \|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|(\delta u,\,\nabla\delta Q)\|_{\dot H^{-\frac12}} \\
&\lesssim \|\nabla Q_2\|^2_{L^2_x}\|\Delta Q_2\|^2_{L^2_x}\|(\delta u,\,\nabla\delta Q)\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}} + C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}.
\end{aligned}\tag{73}$$

When $j = 2$ in (71), we observe that

$$\begin{aligned}
\mathcal{I}^4_2(q,q',q'',q''') := &\int_{\mathbb{R}_2}\mathrm{tr}\Big\{\;\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{(\dot S_{q'''-1}Q_2-\dot S_{q''-1}Q_2)\,\dot\Delta_{q''}\dot\Delta_{q'''}\nabla\delta u\}\,\big]\dot\Delta_q\Delta\delta Q - \\
&\qquad - \dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{(\dot S_{q'''-1}Q_2-\dot S_{q''-1}Q_2)\,\dot\Delta_{q''}\dot\Delta_{q'''}\Delta\delta Q\}\,\big]\dot\Delta_q\nabla\delta u\;\Big\} \\
&\lesssim \|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{(\dot S_{q'''-1}Q_2-\dot S_{q''-1}Q_2)\,\dot\Delta_{q''}\dot\Delta_{q'''}(\nabla\delta u,\,\Delta\delta Q)\,\}\,\big]\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x} \\
&\lesssim 2^q\|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{(\dot S_{q'''-1}Q_2-\dot S_{q''-1}Q_2)\,\dot\Delta_{q''}\dot\Delta_{q'''}(\nabla\delta u,\,\Delta\delta Q)\,\}\,\big]\|_{L^1_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x} \\
&\lesssim 2^q\|\dot\Delta_{q'}Q_2\|_{L^\infty_x}\|(\dot S_{q'''-1}Q_2-\dot S_{q''-1}Q_2)\|_{L^2_x}\|\dot\Delta_{q''}\dot\Delta_{q'''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x} \\
&\lesssim 2^{q-q'-q''}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\|(\dot S_{q'''-1}\nabla Q_2-\dot S_{q''-1}\nabla Q_2)\|_{L^2_x}\|\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x} \\
&\lesssim 2^{q-q'}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\|\nabla Q_2\|_{L^2_x}\|\dot\Delta_{q''}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x},
\end{aligned}$$

which is equivalent to the last inequality of (72) (since $|q''-q'''|\leq 5$). Hence, arguing as for proving (73), the following estimate holds:

$$\sum_{q\in\mathbb{Z}}\sum_{q'\geq q-5}\sum_{q''\leq q'+1}\sum_{|q'''-q''|\leq 5} 2^{-q}\mathcal{I}_2^4(q,q',q'',q''') \lesssim$$
$$\lesssim \|\nabla Q_2\|_{L^2_x}^2\|\Delta Q_2\|_{L^2_x}^2\|(\delta u,\,\nabla\delta Q)\|_{\dot H^{-\frac12}}^2 + C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2 + C_{\Gamma,L}\|\Delta\delta Q\|_{\dot H^{-\frac12}}^2.$$

Now, let us analyze the term in (71) related to $j=3$. Assuming $q''\leq N$ for a suitable positive N, we get

$$\begin{aligned}
\mathcal{I}_3^4(q,q',q'') &:= \int_{\mathbb{R}_2}\mathrm{tr}\big\{\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot S_{q''-1}Q_2\dot\Delta_{q''}\nabla\delta u\}\big]\dot\Delta_q\Delta\delta Q - \dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot S_{q''-1}Q_2\dot\Delta_{q''}\Delta\delta Q\}\big]\dot\Delta_q\nabla\delta u\big\}\\
&\lesssim \|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot S_{q''-1}Q_2\,\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\}\,\big]\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^q\|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot S_{q''-1}Q_2\,\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\}\,\big]\|_{L^1_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^q\|\dot\Delta_{q'}Q_2\|_{L^2_x}\|\dot S_{q''-1}Q_2\|_{L^\infty_x}\|\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^q\|\dot\Delta_{q'}Q_2\|_{L^2_x}(1+\sqrt{N})\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim (1+\sqrt{N})2^{\frac{3q}{2}+\frac{3q''}{2}-2q'}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}2^{-\frac{q''}{2}}\|\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}
\end{aligned}$$

Hence

$$\begin{aligned}
&\sum_{q''\leq N}\sum_{q'\geq q''-1}\sum_{q\leq q'+5}2^{-q}\mathcal{I}_3^4(q,q',q'') \lesssim (1+\sqrt{N})\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|(\nabla\delta u,\Delta\delta Q)\|_{\dot H^{-\frac12}}\times\\
&\qquad\qquad\times\sum_{q''\leq N}2^{-\frac{q''}{2}}\|\dot\Delta_{q''}(\delta u,\nabla\delta Q)\|_{L^2_x}\sum_{q'\geq q''-1}2^{\frac{3q''}{2}-2q'}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\sum_{q\leq q'+5}2^{\frac{q}{2}}\\
&\lesssim (1+\sqrt{N})\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\sum_{q''\leq N}2^{-\frac{q''}{2}}\|\dot\Delta_{q''}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\sum_{q'\geq q''-1}2^{\frac{3q''}{2}-\frac{3q'}{2}}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\\
&\lesssim (1+\sqrt{N})\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|(\delta u,\,\nabla\delta Q)\|_{\dot H^{-\frac12}}\Big(\sum_{q''\in\mathbb{Z}}\Big|\sum_{q'\geq q''-1}2^{\frac32q''-\frac32q'}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\Big|^2\Big)^{\frac12}\\
&\lesssim (1+\sqrt{N})\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|(\delta u,\,\nabla\delta Q)\|_{\dot H^{-\frac12}}.
\end{aligned}$$

Considering the high frequencies $q''>N$

$$\begin{aligned}
\mathcal{I}_3^4(q,q',q'') &\lesssim \|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot S_{q''}Q_2\,\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\}\,\big]\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^q\|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot S_{q''}Q_2\,\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\}\,\big]\|_{L^1_x}\|\dot\Delta_q(\delta u,\,\nabla\delta Q)\|_{L^2_x}\\
&\lesssim 2^q\|\dot\Delta_{q'}Q_2\|_{L^2_x}\|\dot S_{q''}Q_2\|_{L^\infty_x}\|\dot\Delta_{q''}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^{\frac{3q}{2}-2q'}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}(1+\sqrt{q''})\|(Q_2,\,\nabla Q_2)\|_{L^2_x}2^{q''}\|\dot\Delta_{q''}(\delta u,\,\nabla\delta Q)\|_{L^2_x}2^{-\frac{q}{2}}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim (1+\sqrt{q''})2^{\frac{3q}{2}+q''-2q'}\|\Delta Q_2\|_{L^2_x}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|(\delta u,\,\nabla\delta Q)\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}},
\end{aligned}$$

which implies

$$\begin{aligned}
&\sum_{q''>N}\sum_{q'\geq q''-1}\sum_{q\leq q'+5}2^{-q}\mathcal{I}_3^4(q,q',q'') \lesssim\\
&\lesssim \|\Delta Q_2\|_{L^2_x}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|(\delta u,\,\nabla\delta Q)\|_{L^2_x}\sum_{q''>N}(1+\sqrt{q''})2^{q''}\sum_{q'\geq q''-1}2^{-2q'}\sum_{q\leq q'+5}2^{\frac{q}{2}}\\
&\lesssim \|\Delta Q_2\|_{L^2_x}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\|(\delta u,\,\nabla\delta Q)\|_{L^2_x}\sum_{q''>N}(1+\sqrt{q''})2^{q''}\sum_{q'\geq q''-1}2^{-2q'+\frac{q'}{2}}\\
&\lesssim \|\Delta Q_2\|_{L^2_x}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|(\delta u,\,\nabla\delta Q)\|_{L^2_x}\sum_{q''>N}(1+\sqrt{q''})2^{-\frac{q''}{2}}\\
&\lesssim \|\Delta Q_2\|_{L^2_x}\|(Q_2,\,\nabla Q_2)\|_{L^2_x}\|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|(\delta u,\,\nabla\delta Q)\|_{L^2_x}2^{-\frac{N}{2}},
\end{aligned}$$

Summarizing the last inequalities we obtain an estimate similar to (68), so that we can conclude arguing as in (69). Finally, it remains to examine when $j=4$, as last term. Let us define

$$\begin{aligned}
\mathcal{I}_4^4(q,q',q'',q''') := \int_{\mathbb{R}_2}\mathrm{tr}\Big\{\,&\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot\Delta_{q''}(\,\dot\Delta_{q'''}Q_2\,\dot S_{q'''+2}\nabla\delta u\,)\}\,\big]\dot\Delta_q\Delta\delta Q+\\
&-\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot\Delta_{q''}(\,\dot\Delta_{q'''}Q_2\,\dot S_{q'''+2}\Delta\delta Q\,)\}\,\big]\dot\Delta_q\nabla\delta u\,\Big\}
\end{aligned}$$

$$
\begin{aligned}
&\lesssim \|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot\Delta_{q''}[\,\dot\Delta_{q'''}Q_2\,\dot S_{q'''+2}(\nabla\delta u,\,\Delta\delta Q)\,]\,\}\,\big]\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^q\|\dot\Delta_q\big[\,\dot\Delta_{q'}Q_2\mathrm{tr}\{\dot\Delta_{q''}[\,\dot\Delta_{q'''}Q_2\,\dot S_{q'''+2}(\nabla\delta u,\,\Delta\delta Q)\,]\,\}\,\big]\|_{L^1_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^q\|\dot\Delta_{q'}Q_2\|_{L^2_x}\|\dot\Delta_{q''}[\,\dot\Delta_{q'''}Q_2\,\dot S_{q'''+2}(\nabla\delta u,\,\Delta\delta Q)\,]\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^{q+q''}\|\dot\Delta_{q'}Q_2\|_{L^2_x}\|\dot\Delta_{q''}[\,\dot\Delta_{q'''}Q_2\,\dot S_{q'''+2}(\nabla\delta u,\,\Delta\delta Q)\,]\|_{L^1_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^{q+q''}\|\dot\Delta_{q'}\Delta Q_2\|_{L^2_x}\|\dot\Delta_{q'''}Q_2\|_{L^2_x}\|\dot S_{q'''+2}(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}\\
&\lesssim 2^{q-q'+q''-q'''}\|\dot\Delta_{q'}\nabla Q_2\|_{L^2_x}\|\dot\Delta_{q'''}\Delta Q_2\|_{L^2_x}\|\dot S_{q'''+2}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\|\dot\Delta_q(\nabla\delta u,\,\Delta\delta Q)\|_{L^2_x}.
\end{aligned}
$$

Hence, taking the sum in q, q', q'' and q''', we get

$$
\begin{aligned}
&\sum_{q\in\mathbb{Z}}\ \sum_{q'\geq q-5}\ \sum_{q''\leq q'-1}\ \sum_{q'''\geq q''+5} 2^{-q}\mathcal{I}^4_4(q,q',q'',q''')\lesssim\\
&\lesssim \|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|\nabla Q_2\|_{L^2_x}\sum_{q\in\mathbb{Z}}\ \sum_{q'\geq q-5}\ \sum_{q''\leq q'-1}\ \sum_{q'''\geq q''+5} 2^{\frac{q}{2}-q'+q''-q'''}\|\dot\Delta_{q'''}\Delta Q_2\|_{L^2_x}\|\dot S_{q'''+2}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\\
&\lesssim \|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|\nabla Q_2\|_{L^2_x}\sum_{q'''\in\mathbb{Z}}\ \sum_{q''\leq q'''-5} 2^{q''-q'''}\|\dot\Delta_{q'''}\Delta Q_2\|_{L^2_x}\|\dot S_{q'''+2}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\sum_{q'\geq q''+1}2^{-q'}\sum_{q\leq q'+5}2^{\frac{q}{2}}\\
&\lesssim \|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|\nabla Q_2\|_{L^2_x}\sum_{q'''\in\mathbb{Z}}\ \sum_{q''\leq q'''-5} 2^{q''-q'''}\|\dot\Delta_{q'''}\Delta Q_2\|_{L^2_x}\|\dot S_{q'''+2}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\sum_{q'\geq q''+1}2^{-\frac{q'}{2}}\\
&\lesssim \|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|\nabla Q_2\|_{L^2_x}\sum_{q'''\in\mathbb{Z}}\ \sum_{q''\leq q'''-5} 2^{\frac{q''-q'''}{2}}\|\dot\Delta_{q'''}\Delta Q_2\|_{L^2_x}2^{-\frac{q'''}{2}}\|\dot S_{q'''+2}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\\
&\lesssim \|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|\nabla Q_2\|_{L^2_x}\sum_{q'''\in\mathbb{Z}}\|\dot\Delta_{q'''}\Delta Q_2\|_{L^2_x}2^{-\frac{q'''}{2}}\|\dot S_{q'''+2}(\delta u,\,\nabla\delta Q)\|_{L^2_x}\\
&\lesssim \|(\nabla\delta u,\,\Delta\delta Q)\|_{\dot H^{-\frac12}}\|\nabla Q_2\|_{L^2_x}\|\Delta Q_2\|_{L^2_x}\|(\delta u,\,\nabla\delta Q)\|_{\dot H^{-\frac12}}\\
&\lesssim \|\nabla Q_2\|^2_{L^2_x}\|\Delta Q_2\|^2_{L^2_x}\|(\delta u,\,\nabla\delta Q)\|^2_{\dot H^{-\frac12}}+C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}
\end{aligned}
$$

and this concludes the estimates of the term $\mathcal{E}_1+\mathcal{E}_2$.

4.2.5. *Remaining Terms.* For the sake of completeness, now we analyze all the remaining terms. However we point out that they are going to be estimates using simply just Theorem A.1, hence they are not a challenging drawback. For instance, let us observe that

$$
\begin{aligned}
L\langle(\xi\delta D+\delta\Omega)\delta Q,\Delta\delta Q\rangle_{\dot H^{-\frac12}}+L\langle(\xi D_2+\Omega_2)\delta Q,\Delta\delta Q\rangle_{\dot H^{-\frac12}}+L\langle\delta Q(\xi\delta D+\delta\Omega),\Delta\delta Q\rangle_{\dot H^{-\frac12}}+\\
+L\langle\delta Q(\xi D_2+\Omega_2),\Delta\delta Q\rangle_{\dot H^{-\frac12}}\lesssim\|\delta Q\|_{\dot H^{\frac12}}\|\nabla(u_1,\,u_2)\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\lesssim\|\nabla\delta Q\|_{\dot H^{-\frac12}}\times\\
\times\|\nabla(u_1,\,u_2)\|_{L^2_x}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\lesssim\|\nabla(u_1,\,u_2)\|^2_{L^2_x}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}.
\end{aligned}
$$

Moreover $La\Gamma\langle\delta Q,\,\Delta\delta Q\rangle_{\dot H^{-1/2}}\lesssim\|\delta Q\|^2_{\dot H^{-1/2}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-1/2}}$ and

$$
\begin{aligned}
Lb\Gamma\langle Q_1\delta Q+\delta QQ_2,\,\Delta\delta Q\rangle_{\dot H^{-\frac12}}\lesssim\|Q_1\delta Q+\delta QQ_2\|_{\dot H^{-\frac12}}\|\Delta Q\|_{\dot H^{-\frac12}}\lesssim\|(Q_1,\,Q_2)\|_{L^2_x}\|\delta Q\|_{\dot H^{\frac12}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\
\lesssim\|(Q_1,\,Q_2)\|_{L^2_x}\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\lesssim\|(Q_1,\,Q_2)\|^2_{L^2_x}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}.
\end{aligned}
$$

Furthermore, by a direct computation, we get

$$
\begin{aligned}
Lc\Gamma\langle\delta Q\mathrm{tr}\{Q_1^2\},\Delta\delta Q\rangle_{\dot H^{-\frac12}}+Lc\Gamma\langle Q_2\mathrm{tr}\{Q_1\delta Q+\delta QQ_2\},\Delta\delta Q\rangle_{\dot H^{-\frac12}}\lesssim\|(Q_1^2,\,Q_2^2)\|_{L^2_x}\|\delta Q\|_{\dot H^{\frac12}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\
\lesssim\|(Q_1,\,Q_2)\|^2_{L^4_x}\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\lesssim\|(Q_1,\,Q_2)\|^2_{L^2_x}\|\nabla(Q_1,\,Q_2)\|^2_{L^2_x}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}
\end{aligned}
$$

and

$$
\begin{aligned}
L\langle\delta u\cdot\nabla Q_1,\Delta\delta Q\rangle_{\dot H^{-\frac12}}+L\langle u_2\cdot\nabla\delta Q,\,\Delta\delta Q\rangle_{\dot H^{-\frac12}}\lesssim\|(u_2,\nabla Q_1)\|_{\dot H^{\frac34}}\|(\delta u,\,\nabla\delta Q)\|_{\dot H^{-\frac14}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\
\lesssim\|(u_2,\nabla Q_1)\|^{\frac14}_{L^2_x}\|(\nabla u_2,\Delta Q_1)\|^{\frac34}_{L^2_x}\|(\delta u,\,\nabla\delta Q)\|^{\frac34}_{\dot H^{-\frac12}}\|(\Delta\delta u,\,\Delta\delta Q)\|^{\frac14}_{\dot H^{-\frac12}}\|\Delta\delta Q\|_{\dot H^{-\frac12}}\\
\lesssim\|(u_2,\nabla Q_1)\|^{\frac23}_{L^2_x}\|(\nabla u_2,\Delta Q_1)\|^2_{L^2_x}\|(\delta u,\,\nabla\delta Q)\|^2_{\dot H^{-\frac12}}+C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}+C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}}.
\end{aligned}
$$

Moreover $a\xi\langle\delta QQ_1,\,\nabla\delta u\rangle_{\dot H^{-1/2}}\lesssim\|\delta Q\|_{\dot H^{1/2}}\|Q_1\|_{L^2}\|\nabla\delta u\|_{\dot H^{-1/2}}\lesssim\|Q_1\|^2_{L^2}\|\nabla\delta Q\|^2_{\dot H^{-1/2}}+C_\nu\|\nabla\delta u\|^2_{\dot H^{-1/2}}$,

$$
\begin{aligned}
b\xi\langle\delta Q(Q_1^2-\mathrm{tr}\{Q_1^2\}\frac{\mathrm{Id}}{2}),\,\nabla\delta u\rangle_{\dot H^{-\frac12}}\lesssim\|\delta Q\|_{\dot H^{\frac12}}\|Q_1^2\|_{L^2_x}\|\nabla\delta u\|_{\dot H^{-\frac12}}=\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|Q_1\|^2_{L^4_x}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\
\lesssim\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|Q_1\|_{L^2_x}\|\nabla Q_1\|_{L^2_x}\|\nabla\delta u\|_{\dot H^{-\frac12}}\lesssim\|Q_1\|^2_{L^2_x}\|\nabla Q_1\|^2_{L^2_x}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}}+C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}
\end{aligned}
$$

and

$$\begin{aligned}c\xi\langle \delta Q\mathrm{tr}(Q_1^2)Q_1,\, \nabla\delta u\rangle_{\dot H^{-\frac12}} &\lesssim \|\delta Q\|_{\dot H^{\frac12}}\|Q_1^2\|_{L^2_x}\|Q_1\|_{L^\infty}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\lesssim \|Q_1\|^2_{L^2_x}\|\nabla Q_1\|^2_{L^2_x}\|Q_1\|^2_{H^2}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}.\end{aligned}$$

Now, $a\xi\langle (Q_2+\mathrm{Id}/2)\delta Q, \nabla\delta u\rangle_{\dot H^{-1/2}} \lesssim (\|Q_2\|^2_{L^2_x}+1)\|\nabla\delta Q\|^2_{\dot H^{-1/2}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-1/2}}$ and

$$\begin{aligned}&b\xi\langle (Q_2+\frac{\mathrm{Id}}{2})(Q_1\delta Q+\delta QQ_2),\, \nabla\delta u\rangle_{\dot H^{-\frac12}} - b\xi\langle Q_2\mathrm{tr}\{Q_1\delta Q+\delta QQ_2\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} \lesssim\\ &\lesssim (\|Q_2\|_{L^\infty_x}+1)\|(Q_1,\, Q_2)\|_{L^2_x}\|\delta Q\|_{\dot H^{\frac12}}\|\nabla\delta u\|_{\dot H^{-\frac12}} \lesssim (\|Q_2\|_{H^2}+1)\|(Q_1,\, Q_2)\|_{L^2_x}\times\\ &\times\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|\nabla\delta u\|_{\dot H^{-\frac12}} \lesssim (\|Q_2\|_{H^2}+1)^2\|(Q_1,\, Q_2)\|^2_{L^2_x}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}.\end{aligned}$$

Equivalently, we get

$$\begin{aligned}&c\xi\langle (Q_2+\frac{\mathrm{Id}}{2})\delta Q\mathrm{tr}\{Q_1^2\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + c\xi\langle (Q_2+\frac{\mathrm{Id}}{2})Q_2\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} \lesssim\\ &\lesssim \|Q_2\|_{L^\infty}\|(Q_1^2,\, Q_2^2)\|_{L^2_x}\|\delta Q\|_{\dot H^{\frac12}}\|\nabla\delta u\|_{\dot H^{-\frac12}} \lesssim \|Q_2\|_{H^2}\|(Q_1,\, Q_2)\|^2_{L^4_x}\|\nabla\delta Q\|_{\dot H^{\frac12}}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\lesssim \|Q_2\|^2_{H^2}\|(Q_1,\, Q_2)\|^2_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^2_{L^2_x}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}\end{aligned}$$

and moreover

$$\begin{aligned}L\xi\langle \delta Q\Delta\delta Q,\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + L\xi\langle \delta Q\Delta Q_2, \nabla\delta u\rangle_{\dot H^{-\frac12}} &\lesssim \|\delta Q\|_{\dot H^{\frac12}}\|\Delta(Q_1,\, Q_2)\|_{L^2_x}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\lesssim \|\Delta(Q_1,\, Q_2)\|^2_{L^2_x}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}.\end{aligned}$$

We can similarly control the terms from $-a\xi\langle Q_1\delta Q, \nabla\delta u\rangle_{\dot H^{-1/2}}$ to $L\xi\langle \Delta Q_2, \delta Q,\, \nabla\delta\rangle_{\dot H^{-1/2}}$ in (40), proceeding as in the previous estimates. Furthermore

$$\begin{aligned}&2a\xi\langle \delta Q\mathrm{tr}\{Q_1^2\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + 2a\xi\langle Q_2\mathrm{tr}\{\delta QQ_1\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + 2a\xi\langle Q_2\mathrm{tr}\{Q_2\delta Q\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} \lesssim\\ &\lesssim \|\delta Q\|_{\dot H^{\frac12}}\|(Q_1^2,\, Q_2^2)\|_{L^2_x}\|\nabla\delta u\|_{\dot H^{-\frac12}} \lesssim \|(Q_1,\, Q_2)\|^2_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^2_{L^2_x}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}},\end{aligned}$$

$$\begin{aligned}&2b\xi\langle \delta Q\mathrm{tr}\{Q_1^3\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + 2b\xi\langle Q_2\mathrm{tr}\{\delta QQ_2^2\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + 2b\xi\langle Q_2\mathrm{tr}\{Q_2(\delta QQ_1+Q_2\delta Q)\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}}\\ &\qquad\lesssim \|(Q_1,\, Q_2)\|^2_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^2_{L^2_x}\|(Q_1,\, Q_2)\|^2_{H^2}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}\end{aligned}$$

and also

$$\begin{aligned}&2c\xi\langle \delta Q\mathrm{tr}\{Q_1^2\}^2,\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + 2c\xi\langle Q_2\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\}\mathrm{tr}\{Q_1^2\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}}+\\ &+2c\xi\langle Q_2\mathrm{tr}\{Q_2^2\}\mathrm{tr}\{\delta QQ_1+Q_2\delta Q\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} \lesssim \|(Q_1^4,\, Q_2^4)\|_{L^2_x}\|\delta Q\|_{\dot H^{\frac12}}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\lesssim \|(Q_1,\, Q_2)\|^4_{L^8_x}\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|\nabla\delta u\|_{\dot H^{-\frac12}} \lesssim \|(Q_1,\, Q_2)\|_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^3_{L^2_x}\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\lesssim \|(Q_1,\, Q_2)\|^2_{L^2_x}\|\nabla(Q_1,\, Q_2)\|^6_{L^2_x}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}.\end{aligned}$$

Furthermore, we observe that

$$\begin{aligned}&2L\xi\langle \delta Q\mathrm{tr}\{\delta Q\Delta\delta Q\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + 2L\xi\langle \delta Q\mathrm{tr}\{\delta Q\Delta Q_2\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + 2L\xi\langle \delta Q\mathrm{tr}\{Q_2\Delta\delta Q\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}}+\\ &+2L\xi\langle Q_2\mathrm{tr}\{\delta Q\Delta\delta Q\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + 2L\xi\langle \delta Q\mathrm{tr}\{Q_2\Delta Q_2\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + 2L\xi\langle Q_2\mathrm{tr}\{\delta Q\Delta Q_2\},\, \nabla\delta u\rangle_{\dot H^{-\frac12}}\\ &\qquad\lesssim \|\delta Q\|_{\dot H^{\frac12}}\|\Delta(Q_1,\, Q_2)\|_{L^2_x}\|(Q_1,\, Q_2)\|_{L^\infty_x}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\qquad\lesssim \|\Delta(Q_1,\, Q_2)\|^2_{L^2_x}\|(Q_1,\, Q_2)\|^2_{H^2}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}\end{aligned}$$

and

$$\begin{aligned}L\langle \nabla\delta Q\odot\nabla Q_1,\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + L\langle \nabla Q_2\odot\nabla\delta Q,\, \nabla\delta u\rangle_{\dot H^{-\frac12}} &\lesssim \|\nabla\delta Q\|_{\dot H^{-\frac14}}\|\nabla(Q_1,\, Q_2)\|_{\dot H^{\frac34}}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\lesssim \|\nabla\delta Q\|^{\frac34}_{\dot H^{-\frac12}}\|\nabla\delta Q\|^{\frac14}_{\dot H^{\frac12}}\|\nabla(Q_1,\, Q_2)\|^{\frac14}_{L^2}\|\Delta(Q_1,\, Q_2)\|^{\frac34}_{L^2}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\lesssim \|\nabla\delta Q\|^{\frac34}_{\dot H^{-\frac12}}\|\Delta\delta Q\|^{\frac14}_{\dot H^{-\frac12}}\|\nabla(Q_1,\, Q_2)\|^{\frac14}_{L^2}\|\Delta(Q_1,\, Q_2)\|^{\frac34}_{L^2}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\lesssim \|\nabla(Q_1,\, Q_2)\|^{\frac23}_{L^2}\|\Delta(Q_1,\, Q_2)\|^2_{L^2}\|\nabla\delta Q\|^2_{\dot H^{-\frac12}} + C_{\Gamma,L}\|\Delta\delta Q\|^2_{\dot H^{-\frac12}} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}}.\end{aligned}$$

Moreover

$$\begin{aligned}La\langle \delta QQ_1,\, \nabla\delta u\rangle_{\dot H^{-\frac12}} + a\langle Q_2\delta Q,\, \nabla\delta u\rangle_{\dot H^{-\frac12}} - a\langle Q_1\delta Q,\, \nabla\delta u\rangle_{\dot H^{-\frac12}} &\lesssim \|\delta Q\|_{\dot H^{\frac12}}\|(Q_1,\, Q_2)\|_{L^2}\|\nabla\delta u\|_{\dot H^{-\frac12}}\\ &\lesssim \|\nabla\delta Q\|_{\dot H^{-\frac12}}\|(Q_1,\, Q_2)\|_{L^2}\|\nabla\delta u\|_{\dot H^{-\frac12}} \lesssim \|\nabla\delta Q\|^2_{\dot H^{-\frac12}}\|(Q_1,\, Q_2)\|^2_{L^2} + C_\nu\|\nabla\delta u\|^2_{\dot H^{-\frac12}},\end{aligned}$$

$$-Lb\langle\delta Q(Q_1^2-\mathrm{tr}\{Q_1^2\}\frac{\mathrm{Id}}{3}),\,\nabla\delta u\rangle_{\dot H^{-\frac12}}-b\langle Q_2(Q_1\delta Q+\delta QQ_2-\mathrm{tr}\{Q_1\delta Q+\delta QQ_2\}\frac{\mathrm{Id}}{3}),\,\nabla\delta u\rangle_{\dot H^{-\frac12}}+$$

$$+b\langle(Q_1^2-\mathrm{tr}\{Q_1^2\}\frac{\mathrm{Id}}{3})\delta Q,\,\nabla\delta u\rangle_{\dot H^{-\frac12}}-b\langle(Q_1\delta Q+\delta QQ_2-\mathrm{tr}\{Q_1\delta Q+\delta QQ_2\}\frac{\mathrm{Id}}{3})\delta Q,\,\nabla\delta u\rangle_{\dot H^{-\frac12}}\lesssim$$

$$\lesssim\|\delta Q\|_{\dot H^{-\frac12}}\|(Q_1^2,\,Q_2^2)\|_{L^2}\|\nabla\delta u\|_{\dot H^{-\frac12}}\lesssim\|\delta Q\|_{\dot H^{\frac12}}\|(Q_1,\,Q_2)\|_{L^4}^2\|\nabla\delta u\|_{\dot H^{-\frac12}}\lesssim\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|(Q_1,\,Q_2)\|_{L^2}\times$$

$$\times\|\nabla(Q_1,\,Q_2)\|_{L^2}\|\nabla\delta u\|_{\dot H^{-\frac12}}\lesssim\|\nabla\delta Q\|_{\dot H^{-\frac12}}^2\|(Q_1,\,Q_2)\|_{L^2}^2\|\nabla(Q_1,\,Q_2)\|_{L^2}^2+C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2,$$

and

$$Lc\langle\delta QQ_1\mathrm{tr}\{Q_1^2\},\,\nabla\delta u\rangle_{\dot H^{-\frac12}}+c\langle Q_2\delta Q\mathrm{tr}\{Q_1^2\},\,\nabla\delta u\rangle_{\dot H^{-\frac12}}-c\langle Q_1\delta Q\mathrm{tr}\{Q_1^2\},\,\nabla\delta u\rangle_{\dot H^{-\frac12}}-$$

$$-c\langle\delta QQ_2\mathrm{tr}\{Q_1^2\},\,\nabla\delta u\rangle_{\dot H^{-\frac12}}\lesssim\|\delta Q\|_{\dot H^{\frac12}}\|(Q_1,\,Q_2)\|_{L^\infty_x}\|(Q_1^2,\,Q_2^2)\|_{L^2_x}\|\nabla\delta u\|_{\dot H^{-\frac12}}\lesssim\|\nabla\delta Q\|_{\dot H^{-\frac12}}\|(Q_1,\,Q_2)\|_{H^2}\times$$

$$\times\|(Q_1,\,Q_2)\|_{L^4_x}^2\|\nabla\delta u\|_{\dot H^{-\frac12}}\lesssim\|\nabla\delta Q\|_{\dot H^{-\frac12}}^2\|(Q_1,\,Q_2)\|_{H^2}^2\|(Q_1,\,Q_2)\|_{L^2}^2\|\nabla(Q_1,\,Q_2)\|_{L^2}^2+C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2.$$

Finally

$$\langle u_2\cdot\nabla\delta u,\delta u\rangle_{\dot H^{-\frac12}}=-\langle u_2\otimes\delta u,\nabla\delta u\rangle_{\dot H^{-\frac12}}\lesssim\|u_2\|_{\dot H^{\frac12}}\|\delta u\|_{L^2}\|\nabla\delta u\|_{\dot H^{-\frac12}}$$

$$\lesssim\|u_2\|_{L^2}^{\frac12}\|\nabla u_2\|_{L^2}^{\frac12}\|\delta u\|_{\dot H^{-\frac12}}^{\frac12}\|\nabla\delta u\|_{\dot H^{-\frac12}}^{\frac32}\lesssim\|u_2\|_{L^2}^2\|\nabla u_2\|_{L^2}^2\|\delta u\|_{\dot H^{-\frac12}}^2+C_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2$$

and $\langle\delta u\cdot\nabla u_1,\,\delta u\rangle_{\dot H^{-1/2}}\lesssim\|\delta u\|_{\dot H^{1/2}}\|\nabla u_1\|_{L^2}\|\delta u\|_{\dot H^{-1/2}}\lesssim C_\nu\|\nabla\delta u\|_{\dot H^{-1/2}}^2+\|\nabla u_1\|_{L^2}^2\|\delta u\|_{\dot H^{-1/2}}^2$.

4.2.6. *Conclusion.* Recalling (40) and summarizing all the the previous estimates, we conclude that there exists a function χ which belongs to $L^1_{loc}(\mathbb{R}_+)$ such that

$$\frac{\mathrm{d}}{\mathrm{d}t}\Phi(t)+\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2+\Gamma L^2\|\Delta Q\|_{\dot H^{-\frac12}}^2\lesssim\chi(t)\mu(\Phi(t))+c_\nu\|\nabla\delta u\|_{\dot H^{-\frac12}}^2+C_{\Gamma,L}\|\Delta Q\|_{\dot H^{-\frac12}}^2$$

where μ is the Osgood modulus of continuity defined in (42). Hence, choosing $C_{\Gamma,L}$ and C_ν small enough from the beginning, we can absorb the last two terms on the right-hand side by the left-hand side, obtaining (41). We deduce that $\Phi\equiv 0$, thanks to the Osgood Lemma and the null initial data $\Phi(0)=0$. Thus, $(\delta u,\nabla\delta Q)$ is identically zero and δQ as well, since $\delta Q(t)$ decades to 0 at infinity for almost every t.

□

Appendix A.

Theorem A.1. *Let s and t be two real numbers such that $|s|$ and $|t|$ belong to $[0,d/2)$. Let us assume that $s+t$ is positive, then for every $a\in\dot H^s(\mathbb{R}^d)$ and for every $b\in\dot H^t(\mathbb{R}^d)$, the product ab belongs to $\dot H^{s+t-d/2}$ and there exists a positive constant (not dependent by a and b) such that*

$$\|ab\|_{\dot H^{s+t-d/2}}\le C\|a\|_{\dot H^s}\|b\|_{\dot H^t}$$

Proof. At first we identify the Sobolev Spaces $\dot H^s$ and $\dot H^t$ with the Besov Spaces $\dot B^s_{2,2}$ and $\dot B^t_{2,2}$ respectively. We claim that ab belongs to $\dot B^{s+t-d/2}_{2,2}$ and

$$\|ab\|_{\dot B^{s+t-\frac d2}_{2,2}}\le C\|a\|_{\dot B^s_{2,2}}\|b\|_{\dot B^t_{2,2}},$$

for a suitable positive constant.
We decompose the product ab through the Bony decomposition, namely $ab=\dot T_ab+\dot T_ba+R(a,b)$, where

$$\dot T_ab:=\sum_{q\in\mathbb{Z}}\dot\Delta_qa\,\dot S_{q-1}b,\qquad\dot T_ba:=\sum_{q\in\mathbb{Z}}\dot S_{q-1}a\,\dot\Delta_qb,\qquad\dot R(a,b):=\sum_{\substack{q\in\mathbb{Z}\\|\nu|\le1}}\dot\Delta_qa\,\dot\Delta_{q+\nu}b.$$

For any $q\in\mathbb{Z}$, we have

$$2^{q(s+t-\frac d2)}\|(\dot\Delta_q\dot T_ab,\,\dot\Delta_q\dot T_ba)\|_{L^2}\lesssim$$

$$\lesssim\sum_{|q-q'|\le5}2^{q's}\|\dot\Delta_qa\|_{L^2_x}2^{q'(t-\frac d2)}\|\dot S_{q-1}b\|_{L^\infty}+\sum_{|q-q'|\le5}2^{q'(s-\frac d2)}\|\dot S_{q-1}a\|_{L^\infty_x}2^{q't}\|\dot\Delta_qb\|_{L^2},$$

hence

$$\|(\dot T_ab,\,\dot T_ba)\|_{\dot B^{s+t-\frac d2}_{2,2}}\le\|(\dot T_ab,\,\dot T_ba)\|_{\dot B^{s+t-\frac d2}_{2,1}}\lesssim\|a\|_{\dot B^s_{2,2}}\|b\|_{\dot B^{t-\frac d2}_{\infty,2}}+\|a\|_{\dot B^{s-\frac d2}_{\infty,2}}\|b\|_{\dot B^t_{2,2}}\lesssim\|a\|_{\dot B^s_{2,2}}\|b\|_{\dot B^t_{2,2}},$$

where we have used the embedding $\dot B^\sigma_{2,2}\hookrightarrow\dot B^{\sigma-d/2}_{\infty,2}$, for any $\sigma\in\mathbb{R}$ and moreover the following norm-equivalence

$$\|u\|_{\dot B^{\tilde\sigma}_{p,r}}\approx\|(2^{\tilde\sigma}\|S_qu\|_{L^p_x})_{q\in\mathbb{Z}}\|_{l^r(\mathbb{Z})},\qquad u\in\dot B^{\tilde\sigma}_{p,r},$$

for any $1\le p,r\le\infty$ and $\tilde\sigma<0$.

In order to conclude the proof, we have to handle the rest $\dot R(a,b)$. By a direct computation, for any $q\in\mathbb{Z}$,

$$2^{(t+s)q}\|\dot\Delta_q\dot R(a,b)\|_{L^1_x}\le\sum_{\substack{q'\ge q-5\\|\nu|\le1}}2^{(q-q')(s+t)}2^{q's}\|\dot\Delta_{q'}a\|_{L^2_x}2^{(q'+\nu)t}\|\dot\Delta_{q'+\nu}a\|_{L^2_x},$$

so that, thanks to the Young inequality, we deduce

$$\|\dot R(a,b)\|_{\dot B^{s+t-\frac{d}{2}}_{2,2}}\lesssim\|\dot R(a,b)\|_{\dot B^{s+t}_{1,1}}\lesssim\|a\|_{\dot B^{s}_{2,2}}\|b\|_{\dot B^{t}_{2,2}},$$

where we have used the embedding $\dot B^{s+t}_{1,1}\hookrightarrow\dot B^{s+t-d/2}_{2,2}$ and moreover $\sum_{q\le5}2^{q(s+t)}<\infty$ since $s+t$ is positive. $\square$

Theorem A.2. *Let N be a positive real number and f a function in H^1. Then $\dot S_N f$ belongs to L^∞_x and*

$$\|\dot S_N f\|_{L^\infty_x}\lesssim\|f\|_{L^2_x}+\sqrt{N}\|\nabla f\|_{L^2_x}\lesssim(1+\sqrt{N})\|(f,\,\nabla f)\|_{L^2_x}.$$

Proof. We split $\dot S_N f$ into two parts, namely $\dot S_N f=\sum_{q<0}\dot\Delta_q f+\sum_{0\le q<N}\dot\Delta_q f$. First we observe that

$$\|\sum_{q<0}\dot\Delta_q f\|_{L^\infty_x}\le\sum_{q<0}\|\dot\Delta_q f\|_{L^\infty_x}\lesssim\sum_{q<0}2^q\|\dot\Delta_q f\|_{L^2_x}\lesssim\big(\sum_{q<0}2^q\big)\|f\|_{L^2_x}.$$

Similarly, considering the second term, we get

$$\begin{aligned}\|\sum_{0<q\le N}\dot\Delta_q f\|_{L^\infty_x}&\le\sum_{0<q\le N}\|\dot\Delta_q f\|_{L^\infty_x}\lesssim\sum_{0<q\le N}2^q\|\dot\Delta_q f\|_{L^2_x}\\&\lesssim\sum_{0<q\le N}\|\dot\Delta_q\nabla f\|_{L^2_x}\lesssim\Big(\sum_{0<q\le N}1\Big)^{\frac12}\Big(\sum_{0<q\le N}\|\dot\Delta_q\nabla f\|^2_{L^2_x}\Big)^{\frac12}\lesssim\sqrt{N}\|f\|_{\dot H^1},\end{aligned}$$

which concludes the proof of the Theorem. $\square$

The following Lemma plays a main role in the uniqueness result of Theorem 1.1, more precisely inequality (74) is the key for the double-logarithmic estimate.

Lemma A.3. *There exist a positive constant C such that for any $p\in[1,\infty)$ the following inequality is satisfied:*

$$\|f\|_{L^{2p}(\mathbb{R}^2)}\le C\sqrt{p}\|f\|^{\frac1p}_{L^2(\mathbb{R}^2)}\|\nabla f\|^{1-\frac1p}_{L^2(\mathbb{R}^2)}\tag{74}$$

Proof. The proof of this lemma was presented in [32] (lemma 4.3) and we report it here, for the sake of simplicity. thanks to Sobolev embeddings, we have

$$\|f\|_{L^{2p}(\mathbb{R}^2)}\le C\sqrt{p}\|f\|_{\dot H^{1-\frac1p}(\mathbb{R}^2)}.\tag{75}$$

Moreover, since $\dot H^{1-1/p}(\mathbb{R}^2)$ is an interpolation space between $L^2(\mathbb{R}^2)$ and $\dot H^1(\mathbb{R}^2)$, the following inequality is satisfied:

$$\|f\|_{\dot H^{1-\frac1p}(\mathbb{R}^2)}\le\|f\|^{\frac1p}_{L^2(\mathbb{R}^2)}\|\nabla f\|^{1-\frac1p}_{L^2(\mathbb{R}^2)},$$

which leads to (74), together with (75). $\square$

Appendix B.

Proposition B.1. *Let $(Q^{(n)},u^n)$ be a smooth solution of (27) in dimension $d=2$ or $d=3$, with restriction (3), and smooth initial data $(\bar Q(x),\bar u(x))$, that decays fast enough at infinity so that we can integrate by parts in space (for any $t\ge0$) without boundary terms. We assume that $|\xi|<\xi_0$ where ξ_0 is an explicitly computable constant, scale invariant, depending on $a,b,c,d,\Gamma,\nu,\lambda$.*

For $(\bar Q,\bar u)\in H^1\times L^2$,we have

$$\|Q^{(n)}(t,\cdot)\|_{H^1}\le C_1+\bar C_1e^{\bar C_1t}\|\bar Q\|_{H^1},\forall t\ge0\tag{76}$$

with $C_1,\bar C_1$ depending on $(a,b,c,d,\Gamma,L,\nu,\bar Q,\bar u)$. Moreover

$$\|u^n(t,\cdot)\|^2_{L^2}+\nu\int_0^t\|\nabla u^n\|^2_{L^2}\le C_1\tag{77}$$

Proof. We denote:

$$X^n_{\alpha\beta} \overset{\text{def}}{=} L\Delta Q^{(n)}_{\alpha\beta} - cQ^{(n)}_{\alpha\beta}\mathrm{tr}((Q^{(n)})^2),\ \alpha,\beta = 1,2,3. \tag{78}$$

Multiplying the first equation in (27) by $-\lambda\bar{H}^n$ and the second one by u^n,taking the trace and integrating over $\mathbb{R}^d$, we get

$$\begin{aligned}
&\frac{d}{dt}\int_{\mathbb{R}^d}\frac{1}{2}|u^n|^2 + \frac{L\lambda}{2}|\nabla Q^{(n)}|^2 + \lambda\big(\frac{a}{2}|Q^{(n)}|^2 - \frac{b}{3}\mathrm{tr}(Q^{(n)})^3 + \frac{c}{4}|Q^{(n)}|^4\big)\,dx + \nu\|\nabla u^n\|^2_{L^2} + \Gamma\lambda L^2\|\Delta Q^{(n)}\|^2_{L^2}\\
&+\Gamma\lambda c^2\|J_n(Q^{(n)}\mathrm{tr}\{Q^{(n)}\})\|^2_{L^2} - 2cL\Gamma\lambda\int_{\mathbb{R}^d}\Delta Q^{(n)}_{\alpha\beta}Q^{(n)}_{\alpha\beta}\mathrm{tr}\{(Q^{(n)})^2\}\,dx + a^2\Gamma\lambda\|Q^{(n)}\|^2_{L^2}\\
&+b^2\Gamma\lambda\int_{\mathbb{R}^d}\mathrm{tr}\big\{J_n\big((Q^{(n)})^2 - \frac{\mathrm{tr}\{(Q^{(n)})^2\}}{d}\big)^2\big\}\,dx + \varepsilon\int_{\mathbb{R}^d}|R_\varepsilon u\nabla Q^{(n)}|^3\,dx + \varepsilon\int_{\mathbb{R}^d}|R_\varepsilon\nabla u^n|^4\,dx\\
&\leq 2a\Gamma\lambda\underbrace{\int_{\mathbb{R}^d}\mathrm{tr}\{X^nQ^{(n)}\}\,dx}_{\overset{\text{def}}{=}\mathcal{I}_n} - 2b\Gamma\lambda\underbrace{\int_{\mathbb{R}^d}\mathrm{tr}\{X^n(Q^{(n)})^2\}\,dx}_{\overset{\text{def}}{=}\mathcal{J}_n}\\
&+2ab\Gamma\lambda\int_{\mathbb{R}^d}\mathrm{tr}\{(Q^{(n)})^3\}\,dx + \lambda\underbrace{\int_{\mathbb{R}^d}J_n\left(R_\varepsilon u^n\cdot\nabla Q^{(n)}_{\alpha\beta}\right)J_n\left(bQ^{(n)}_{\alpha\gamma}Q^{(n)}_{\gamma\beta} - cQ^{(n)}_{\alpha\beta}\big|Q^{(n)}\big|^2\right)dx}_{\overset{\text{def}}{=}\mathcal{II}}\\
&+\lambda\int_{\mathbb{R}^d}J_n\left(-R_\varepsilon\Omega^{(n)}_{\alpha\gamma}Q^{(n)}_{\gamma\beta} + Q^{(n)}_{\alpha\gamma}R_\varepsilon\Omega^{(n)}_{\gamma\beta}\right)J_n\left(bQ^{(n)}_{\alpha\delta}Q^{(n)}_{\delta\beta} - cQ^{(n)}_{\alpha\beta}\big|Q^{(n)}\big|^2\right)dx
\end{aligned} \tag{79}$$

Integrating by parts we have:

$$\begin{aligned}
-2cL\Gamma\lambda\int_{\mathbb{R}^d}\Delta Q^{(n)}_{\alpha\beta}Q^{(n)}_{\alpha\beta}\mathrm{tr}\{(Q^{(n)})^2\}dx &= 2cL\Gamma\lambda\int_{\mathbb{R}^d}Q^{(n)}_{\alpha\beta,k}Q^{(n)}_{\alpha\beta,k}\mathrm{tr}\{(Q^{(n)})^2\}dx\\
+2cL\Gamma\lambda\int_{\mathbb{R}^d}Q^{(n)}_{\alpha\beta,k}Q^{(n)}_{\alpha\beta}\partial_k\left(\mathrm{tr}\{(Q^{(n)})^2\}\right)dx &= 2cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla Q^{(n)}|^2\mathrm{tr}\{(Q^{(n)})^2\})\,dx\\
&+cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla\left(\mathrm{tr}\{(Q^{(n)})^2\}\right)|^2\,dx \geq 0
\end{aligned} \tag{80}$$

(where for the last inequality we used the assumption (3) and $L,\Gamma,\lambda > 0$). One can easily see that

$$\mathcal{I}_n = -\frac{L}{2}\|\nabla Q^{(n)}\|^2_{L^2} - c\|Q^{(n)}\|^4_{L^4} \tag{81}$$

and moreover

$$\begin{aligned}
&\lambda\int_{\mathbb{R}^d}J_n\left(-R_\varepsilon\Omega^{(n)}_{\alpha\gamma}Q^{(n)}_{\gamma\beta} + Q^{(n)}_{\alpha\gamma}R_\varepsilon\Omega^{(n)}_{\gamma\beta}\right)J_n\left(bQ^{(n)}_{\alpha\delta}Q^{(n)}_{\delta\beta} - cQ^{(n)}_{\alpha\beta}\big|Q^{(n)}\big|^2\right)dx \leq\\
&\leq\frac{\varepsilon}{2}\int_{\mathbb{R}^d}|R_\varepsilon\nabla u^n|^4\,dx + C(\varepsilon)\int_{\mathbb{R}^d}|Q^{(n)}|^4\,dx + \frac{\Gamma c^2}{2}\int_{\mathbb{R}^d}|J_n(Q^{(n)}|Q^{(n)}|^2)|^2\,dx
\end{aligned}$$

On the other hand, for any $\varepsilon > 0$ and $\tilde{C} = \tilde{C}(\varepsilon, c)$ an explicitly computable constant, we have:

$$\begin{aligned}
\mathcal{J}_n &= L\int_{\mathbb{R}^d}Q^{(n)}_{\alpha\beta,kk}Q^{(n)}_{\alpha\gamma}Q^{(n)}_{\gamma\beta}\,dx - c\int_{\mathbb{R}^d}\mathrm{tr}\{(Q^{(n)})^2\}\mathrm{tr}\{(Q)^{(n)})^3\}\,dx \leq -L\int_{\mathbb{R}^d}Q^{(n)}_{\alpha\beta,k}Q^{(n)}_{\alpha\gamma,k}Q^{(n)}_{\gamma\beta}\,dx\\
&-L\int_{\mathbb{R}^d}Q^{(n)}_{\alpha\beta,k}Q^{(n)}_{\alpha\gamma}Q^{(n)}_{\gamma\beta,k} + \int_{\mathbb{R}^d}\mathrm{tr}\{(Q^{(n)})^2\}\left(\frac{\tilde{C}}{\varepsilon}\mathrm{tr}\{(Q^{(n)})^2\} + \varepsilon\mathrm{tr}^2\{(Q^{(n)})^2\}\right)dx\\
&\leq L\varepsilon\int_{\mathbb{R}^d}|\nabla Q^{(n)}|^2\mathrm{tr}\{(Q^{(n)})^2\}\,dx + \frac{\tilde{C}}{\varepsilon}\|\nabla Q^{(n)}\|^2_{L^2} + \int_{\mathbb{R}^d}\mathrm{tr}\{(Q^{(n)})^2\}\left(\frac{\tilde{C}}{\varepsilon}\mathrm{tr}\{(Q^{(n)})^2\} + \varepsilon\mathrm{tr}^2\{(Q^{(n)})^2\}\right)dx
\end{aligned} \tag{82}$$

Using the last four relations in (79) and considering (29) we obtain:

$$\begin{aligned}
&\frac{d}{dt}\int_{\mathbb{R}^d}\frac{1}{2}|u^n|^2 + \frac{L\lambda}{2}|\nabla Q^{(n)}|^2 + \lambda\big(\frac{a}{2}|Q^{(n)}|^2 - \frac{b}{3}\mathrm{tr}(Q^{(n)})^3 + \frac{c}{4}|Q^{(n)}|^4\big)\,dx + \nu\|\nabla u^n\|^2_{L^2} + \Gamma\lambda L^2\|\Delta Q^{(n)}\|^2_{L^2}\\
&+\frac{\Gamma\lambda c^2}{2}\|J_n(Q^{(n)}\mathrm{tr}\{Q^{(n)}\})\|^2_{L^2} + a^2\Gamma\lambda\|Q^{(n)}\|^2_{L^2} + 2cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla Q^{(n)}|^2\mathrm{tr}\{(Q^{(n)})^2\}\,dx\\
&+cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla\left(\mathrm{tr}\{(Q^{(n)})^2\}\right)|^2\,dx + \frac{\varepsilon}{2}\int_{\mathbb{R}^d}|R_\varepsilon u^n\cdot\nabla Q^{(n)}|^3\,dx + \frac{\varepsilon}{2}\int_{\mathbb{R}^d}|\nabla R_\varepsilon u^n|^4\,dx\\
&\leq 2|a|\Gamma\lambda(\frac{L}{2}\|\nabla Q^{(n)}\|^2_{L^2} + c\|Q^{(n)}\|^4_{L^4}) + 2|b|\Gamma\lambda L\varepsilon\int_{\mathbb{R}^d}|\nabla Q^{(n)}|^2\mathrm{tr}\{(Q^{(n)})^2\}\,dx + 2|b|\Gamma\lambda\frac{\tilde{C}}{\varepsilon}\|\nabla Q^{(n)}\|^2_{L^2}
\end{aligned}$$

$$+2|b|\Gamma\lambda\int_{\mathbb{R}^d}\mathrm{tr}\{(Q^{(n)})^2\}\left(\frac{\tilde{C}}{\varepsilon}\mathrm{tr}\{(Q^{(n)})^2\}+\varepsilon\mathrm{tr}^2\{(Q^{(n)})^2\}\right)dx+2|ab|\Gamma\lambda(\varepsilon\|Q^{(n)}\|_{L^2}^2+(C(\varepsilon)+\frac{\tilde{C}}{\varepsilon})\|Q^{(n)}\|_{L^4}^4)$$

Taking ε small enough we can absorb all the terms with an ϵ coefficient on the right into the left hand side, and we are left with

$$\frac{d}{dt}\int_{\mathbb{R}^d}\frac{1}{2}|u^n|^2+\frac{L\lambda}{2}|\nabla Q^{(n)}|^2+\lambda\big(\frac{a}{2}|Q^{(n)}|^2-\frac{b}{3}\mathrm{tr}(Q^{(n)})^3+\frac{c}{4}|Q^{(n)}|^4\big)\,dx$$
$$+\nu\|\nabla u^n\|_{L^2}^2+\Gamma\lambda L^2\|\Delta Q^{(n)}\|_{L^2}^2+\frac{\Gamma\lambda c^2}{2}\|J_n(Q^{(n)}\mathrm{tr}\{Q^{(n)})\})\|_{L^2}^2+\Gamma\lambda a^2\|Q^{(n)}\|_{L^2}^2$$
$$+2cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla Q^{(n)}|^2\mathrm{tr}\{(Q^{(n)})^2\}\,dx+cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla\left(\mathrm{tr}\{(Q^{(n)})^2\}\right)|^2\,dx\le\bar{C}\left(\|\nabla Q^{(n)}\|_{L^2}^2+\|Q^{(n)}\|_{L^4}^4\right)$$

with $\bar{C}=\bar{C}(a,b,c)$.

The last relation is not yet enough because there are no positive terms. However, let us note that, if $a>0$ we obtain the a-priori estimates by using the inequality $tr\{(Q^{(n)})^3\}\le\frac{3}{8}tr\{(Q^{(n)})^2)\}+tr\{(Q^{(n)})^2\}^2$. If $a\le 0$ we have to estimate separately $\|Q^{(n)}\|_{L^2}$ and this ask for a smallness condition for ξ. Indeed, proceeding as for proving (25), we get

$$\frac{d}{dt}\Big[\int_{\mathbb{R}^d}\frac{1}{2}|u^n|^2+\frac{L\lambda}{2}|\nabla Q^{(n)}|^2+\lambda\big(\frac{a}{2}|Q^{(n)}|^2-\frac{b}{3}\mathrm{tr}(Q^{(n)})^3+\frac{c}{4}|Q^{(n)}|^4\big)\,dx+M\|Q\|_{L^2}^2\Big]$$
$$+\nu\|\nabla u^n\|_{L^2}^2+\Gamma\lambda L^2\|\Delta Q^{(n)}\|_{L^2}^2+\frac{\Gamma\lambda c^2}{2}\|J_n(Q^{(n)}\mathrm{tr}\{Q^{(n)})\})\|_{L^2}^2+a^2\|Q^{(n)}\|_{L^2}^2$$
$$+2cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla Q^{(n)}|^2\mathrm{tr}\{(Q^{(n)})^2\}\,dx+cL\Gamma\lambda\int_{\mathbb{R}^d}|\nabla\left(\mathrm{tr}\{(Q^{(n)})^2\}\right)|^2\,dx$$
$$+\frac{\varepsilon}{2}\int_{\mathbb{R}^d}|R_\varepsilon u^n\cdot\nabla Q^{(n)}|^3\,dx+\frac{\varepsilon}{2}\int_{\mathbb{R}^d}|\nabla R_\varepsilon u^n|^4\,dx\le\bar{C}\left(\|\nabla Q^{(n)}\|_{L^2}^2+\|Q^{(n)}\|_{L^4}^4\right)+$$
$$MC(d)\varepsilon\int_{\mathbb{R}^d}|\nabla u^n|^2\,dx+\frac{M|\xi|^2}{\varepsilon}\int_{\mathbb{R}^d}|Q^{(n)}|^2+|J_n(Q^{(n)}\mathrm{tr}\{(Q^{(n)})^2\}|^2\,dx+M\hat{C}\int_{\mathbb{R}^d}|Q^{(n)}|^2+|Q^{(n)}|^4\,dx \tag{83}$$

We chose ε small enough so that $MC(d)\varepsilon<\nu$. Finally we make the assumption that $|\xi|$ is small enough, depending on a,b,c,d,ν so that

$$\frac{M|\xi|^2}{\varepsilon}\le\Gamma\lambda c^2$$

Then taking into account that

$$\frac{M}{2}\mathrm{tr}\{(Q^{(n)})^2\}+\frac{c}{8}\mathrm{tr}^2\{(Q^{(n)})^2\}\le(M+\frac{a}{2})\mathrm{tr}\{(Q^{(n)})^2\}-\frac{b}{3}\mathrm{tr}\{(Q^{(n)})^3\}+\frac{c}{4}\mathrm{tr}^2\{(Q^{(n)})^2\}$$

we obtain the claimed relation (76).

□

References

[1] H. Abels, G. Dolzmann and Y. Liu, Well-posedness of a fully coupled Navier-Stokes/Q-tensor system with inhomogeneous boundary data. SIAM J. Math. Anal. 46 (2014), no. 4, 3050-3077

[2] H. Bahouri, J. Chemin, and R. Danchin, Fourier analysis and nonlinear partial differential equations, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 343, Springer, Heidelberg, 2011. MR 2768550 (2011m:35004)

[3] J. W. Barrett, C. Schwab and E. Süli, Existence of global weak solutions for some polymeric flow models. Math. Models Methods Appl. Sci. 15 (2005), no. 6, 939–983

[4] J.-M. Bony. Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires. Ann. Sci. École Norm. Sup. (4), 14(2):209–246, 1981.

[5] A. N. Beris, B. J. Edwards, Thermodynamics of Flowing Systems with Internal Microstructure. Oxford Engineering Science Series, vol. 36, Oxford University Press, Oxford, New York, 1994.

[6] H. Brezis and T. Gallouet, Nonlinear Schrödinger evolution equations. Nonlinear Anal. 4 (1980), no. 4, 677–681

[7] J.Y. Chemin and C.J. Xu, Inclusions de Sobolev en calcul de Weyl-Hörmander et champs de vecteurs sous-elliptiques. (French) [Sobolev embeddings in Weyl-Hörmander calculus and subelliptic vector fields] Ann. Sci. École Norm. Sup. (4) 30 (1997), no. 6, 719–751.

[8] P. Constantin, Nonlinear Fokker-Planck Navier-Stokes systems. Commun. Math. Sci. 3 (2005), no. 4, 531–544

[9] P. Constantin, C. Fefferman, E.S. Titi and A. Zarnescu, Regularity of coupled two-dimensional nonlinear Fokker-Planck and Navier-Stokes systems. Comm. Math. Phys. 270 (2007), no. 3, 789–811.

[10] P. Constantin and N. Masmoudi, Global well-posedness for a Smoluchowski equation coupled with Navier-Stokes equations in 2D. Comm. Math. Phys. 278 (2008), no. 1, 179–191.

[11] P. Constantin and G. Seregin, Global regularity of solutions of coupled Navier-Stokes equations and nonlinear Fokker-Planck equations, arXiv:0901.4462

[12] J. Y. Chemin, Perfect incompressible fluids. Oxford Lecture Series in Mathematics and its Applications, 14. The Clarendon Press, Oxford University Press, New York, 1998

[13] J.Y. Chemin, and N. Masmoudi, About lifespan of regular solutions of equations related to viscoelastic fluids. SIAM J. Math. Anal. 33 (2001), no. 1, 84–112

[14] C.Denniston, E. Orlandini, E; J.M. Yeomans, Lattice Boltzmann simulations of liquid crystals hydrodynamics, Phys. Rev. E., vol. 63, (5), 056702

[15] G. Furioli, P. Lemarié-Rieusset, and E. Terraneo, Unicité dans $L^3(\mathbb{R}^3)$ et dautres espaces fonctionnels limites pour Navier-Stokes, Rev. Mat. Iberoamericana **16** (2000), no. 3, 605–667. MR 1813331 (2002j:76036)

[16] F.Guillén-González, M.A. Rodríguez-Bellido, Weak solutions for an initial-boundary Q-tensor problem related to liquid crystals, Nonlinear Anal., 112, 2015, 84–104

[17] F. Guillén-González, and M.A. Rodríguez-Bellido, Weak time regularity and uniqueness for a Q-tensor model. SIAM Journal on Mathematical Analysis 46, no. 5 (2014): 3540-3567.

[18] J. Huang, S. Ding, Global well-posedness for the dynamical Q -tensor model of liquid crystals, Science China Mathematics June 2015, Volume 58, Issue 6, pp 1349-1366

[19] F. M. Leslie, Some constitutive equations for liquid crystals, Arch. Rat. Mech. Anal. 28, 265 (1968)

[20] J. Li, E. S. Titi, and Z. Xin, On the uniqueness of weak solutions to the Ericksen-Leslie liquid crystal model in $\mathbb{R}^2$, arXiv:1410.1119

[21] F. H. Lin and C.Liu, Nonparabolic dissipative systems modeling the flow of liquid crystals. Comm. Pure Appl. Math. 48 (1995), no. 5, 501–537.

[22] F. H. Lin and C. Liu, Existence of solutions for the Ericksen-Leslie system. Arch. Ration. Mech. Anal. 154 (2000), no. 2, 135–156.

[23] P. G. De Gennes, The physics of liquid crystals. Oxford, Clarendon Press. 1974.

[24] F. H. Lin and C. Liu, Static and Dynamic Theories of Liquid Crystals. Journal of Partial Differential Equations, 14, no. 4, 289–330 (2001).

[25] P.L. Lions and N. Masmoudi, Global solutions for some Oldroyd models of non-Newtonian flows. Chinese Ann. Math. Ser. B 21 (2000), no. 2, 131–146.

[26] F. H. Lin, Fanghua, P. Zhang and Z. Zhang, On the global existence of smooth solution to the 2-D FENE dumbbell model. Comm. Math. Phys. 277 (2008), no. 2, 531–553.

[27] F. H. Lin, J. Y. Lin and C. Wang, Liquid crystal flows in two dimensions. Arch. Rat. Mech. Anal., in press

[28] A. Majumdar and A. Zarnescu, Landau-de Gennes theory of nematic liquid crystals: the Oseen-Frank limit and beyond, Arch. Rat. Mech. Anal., 196 (2010), no.1, 227-280

[29] F. Marchand and M. Paicu, Remarques sur l'unicité pour le système de Navier-Stokes tridimensionnel, C. R. Math. Acad. Sci. Paris **344** (2007), no. 6, 363–366. MR 2309504 (2007m:35189)

[30] N. Masmoudi, Well-posedness for the FENE dumbbell model of polymeric flows. Comm. Pure Appl. Math. 61 (2008), no. 12, 1685–1714

[31] M. Paicu, Équation periodique de Navier-Stokes sans viscosité dans une direction. Comm. Partial Differential Equations 30 (2005), no. 7-9, 1107–1140.

[32] M. Paicu, Équation anisotrope de Navier-Stokes dans des espaces critiques, Rev. Mat. Iberoamericana **21** (2005), no. 1, 179–235. MR 2155019 (2006f:35210)

[33] A. M. Sonnet, P.L. Maffettone, E.G. Virga, Continuum theory for nematic liquid crystals with tensorial order. J. Non-Newtonian Fluid Mech. 119 (2004), 51–59.

[34] G. Toth, C. Denniston and J.M. Yeomans, Hydrodynamics of domain growth in nematic liquid crystals, Phys. Rev. E, 67, 051705 (2003)

[35] Marius Paicu and Arghir Zarnescu, Energy dissipation and regularity for a coupled Navier-Stokes and Q-tensor system, Arch. Ration. Mech. Anal. 203 (2012), no. 1, 45–67. MR 2864407 (2012m:76012)

[36] M. Paicu and A. Zarnescu, Global existence and regularity for the full coupled Navier-Stokes and Q-tensor system , SIAM J. Math. Anal., 43 (2011), no.5, 2009-2049

[37] M.E. Schonbek, Existence and decay of polymeric flows. SIAM J. Math. Anal. 41 (2009), no. 2, 564–587

[38] M. Wang, W. Wang and Z. Zhang, On the uniqueness of weak solution for the 2-D Ericksen–Leslie system, Discrete and Continuous Dynamical Systems - Series B 21 (2016), no. 3, 919–941